Algebraic Topology Exam Questions

This is a collection of questions taken from Algebraic Topology exams over a number of years. To some extent, they have been modified to be compatible with the current version of the course, but some differences remain.

1 Compactness and the Hausdorff property

Some questions in this section use ideas that are specific to metric spaces rather than general topological spaces. These ideas are not developed in the current version of the course.

(1) Let X be a metric space.

- (a) Let Y be a compact subspace of X. Prove that Y is closed in X.
- (b) Let Y and Z be two compact subspaces of X. Prove that $Y \cup Z$ is compact.
- (c) Deduce (or prove otherwise) that every finite space is compact.
- (d) Let Y and Z be compact metric spaces. Prove that $Y \times Z$ is compact.
- (e) Conversely, let Y and Z be metric spaces such that $Z \neq \emptyset$ and $Y \times Z$ is compact. Prove that Y is compact.
- (f) Put $X = \{(x, y, z) \in \mathbb{R}^3 : x^4 + y^4 + z^4 = 1\}$. Prove that X is compact. You may use general theorems provided that you state them precisely.
- (2) Let X be a metric space.
 - (a) Let Y be a compact subspace of X. Prove that Y is closed in X.
 - (b) Let Y and Z be two compact subspaces of X. Prove that $Y \cup Z$ is compact.
 - (c) Deduce (or prove otherwise) that every finite space is compact.
 - (d) Let Y and Z be compact metric spaces. Prove that $Y \times Z$ is compact.
 - (e) Conversely, let Y and Z be metric spaces such that $Z \neq \emptyset$ and $Y \times Z$ is compact. Prove that Y is compact.
 - (f) Put $X = \{(x, y, z) \in \mathbb{R}^3 : x^4 + y^4 + z^4 = 1\}$. Prove that X is compact. You may use general theorems provided that you state them precisely.

(3)

- (a) What does it mean to say that a metric space X is *compact*? (3 marks)
- (b) Let $f: X \to Y$ be a continuous surjective map of metric spaces, where X is compact. Prove that Y is compact. (6 marks)
- (c) Let Z be a closed subset of a compact space X. Prove that Z is compact. (6 marks)

(d) Put $U = \{z \in \mathbb{C} \mid 0 \leq \operatorname{Re}(z) \leq 1\}$, and define $g \colon \mathbb{C} \to \mathbb{C}$ by $g(z) = e^z$.

- (i) Is U compact? (2 marks)
- (ii) Is g(U) compact? (4 marks)
- (iii) Is g(g(U)) compact? (4 marks)

Justify your answers.

- (a) What does it mean to say that a metric space X is compact? (3 marks)
- (b) Let X and Y be compact metric spaces. Prove that $X \times Y$ is compact. (8 marks)
- (c) Let $f: I \to Y$ be a continuous map (where I = [0, 1]). Prove that f(I) is closed in Y. (7 marks)
- (d) Put $X = \mathbb{Z} \times \mathbb{Z}$ and $Y = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 < 4\}$, considered as subspaces of the plane \mathbb{R}^2 .
 - (i) Is X compact? (2 marks)
 - (ii) Is Y compact? (2 marks)
 - (iii) Is $X \cap Y$ compact? (3 marks)

Justify your answers.

(5) 2021-22 Q2:

- (a) Define what is meant by a *topology* on a set X. (3 marks)
- (b) What does it mean to say that a topological space X is *Hausdorff*?(If your definition relies on any other concepts, then you should define them.) (3 marks)
- (c) What does it mean to say that a topological space X is *compact*?(If your definition relies on any other concepts, then you should define them.) (3 marks)
- (d) Let X and Y be topological spaces, and let $f: X \to Y$ be a continuous injective map. For each of the claims below, give a proof or a counterexample with justification.
 - (i) If X is Hausdorff, then Y must also be Hausdorff. (4 marks)
 - (ii) If X is compact, then Y must also be compact. (4 marks)
 - (iii) If Y is Hausdorff, then X must also be Hausdorff. (4 marks)
 - (iv) If Y is compact, then X must also be compact. (4 marks)

(6) 2022-23 Q2:

- (a) What does it mean to say that a topological space X is *compact*? If your explanation relies on any auxiliary terms, then you should define them. (3 marks)
- (b) Let X be compact topological space, and let Y be a closed subset of X.
 - (i) Define the subspace topology on Y. (2 marks)
 - (ii) Prove that when equipped with the subspace topology, Y is again compact. (5 marks)
 - (iii) Give an example of a compact space X and a compact subpace Y such that Y is not closed in X. (3 marks)
 - (iv) Explain a commonly-satisfied condition on X that guarantees that compact subspaces are closed. If your explanation relies on any auxiliary terms, then you should define them. However, you need not prove anything. (3 marks)
- (c) Put $X = \mathbb{Z} \times \mathbb{Z}$ and $Y = \{(x, y) \in \mathbb{R}^2 \mid 100 < x^2 + y^2 < 10000\}$, considered as subspaces of the plane \mathbb{R}^2 .
 - (i) Is X compact? (1 marks)
 - (ii) Is Y compact? (1 marks)
 - (iii) Is $X \cap Y$ compact? (2 marks)

Justify your answers.

(d) Let X be a metric space such that $X \setminus \{x\}$ is compact for all $x \in X$. Prove that X is finite. (5 marks)

(7) 2023-24 Q1:

- (a) What does it mean to say that a topological space X is *Hausdorff*?(If your definition relies on any other concepts, then you should define them.) (3 marks)
- (b) What does it mean to say that a topological space X is compact?(If your definition relies on any other concepts, then you should define them.) (3 marks)
- (c) Put $X = \{(x, y, z) \in \mathbb{R}^3 : x^4 + y^4 + z^4 = 1\}$. Prove that X is compact. You may use general theorems provided that you state them precisely. (5 marks)
- (d) Put $Y = \{(x, y, z) \in \mathbb{R}^3 : x^4 + y^4 + z^4 < 1\}$. Prove that Y is not compact. Here you should argue directly from the definitions and not use any theorems. (5 marks)
- (e) Let Y and Z be two compact subspaces of a topological space X. Prove that $Y \cup Z$ is also compact. (4 marks)
- (f) Let Y and Z be topological spaces such that $Z \neq \emptyset$ and $Y \times Z$ is compact. Prove that Y is compact. You may use standard results so long as you state them clearly and verify carefully that they are applicable. (5 marks)

2 Path components

(8)

- (a) What does it mean to say that a topological space X is *path-connected*?
- (b) Prove that the space S^n is path-connected for all n > 0.
- (c) Let X be a subset of \mathbb{R}^n , and let a be a point in X. What does it mean to say that X is *star-shaped* around a? Show that if X is star-shaped around a, then it is path-connected.
- (d) Suppose that $f: X \to \mathbb{R}$ is continuous, f(x) is nonzero for all x, and there exist $x_0, x_1 \in X$ with $f(x_0) < 0 < f(x_1)$. Prove that X is not path-connected.
- (e) Recall that $GL_3(\mathbb{R})$ is the space of 3×3 invertible matrices over \mathbb{R} . Prove that this space is not path-connected.

(9)

- (a) Let X be a topological space. Define the equivalence relation ~ on X such that $\pi_0(X) = X/\sim$, and prove that it is an equivalence relation.
- (b) Let $f: X \to Y$ be a continuous map. Define the induced map $f_*: \pi_0(X) \to \pi_0(Y)$, and prove that it is well-defined.
- (c) Show that if $f, g: X \to Y$ are homotopic maps then $f_* = g_*: \pi_0(X) \to \pi_0(Y)$.
- (d) Put $X = [-3, -2] \cup [-1, 1] \cup [2, 3]$ and $Y = [0, 1] \cup [2, 10]$, and define $f: X \to Y$ by $f(x) = x^2$. Describe the sets $\pi_0(X)$ and $\pi_0(Y)$ and the map $f_*: \pi_0(X) \to \pi_0(Y)$.

(10)

- (a) Let X be a topoological space. Define the equivalence relation ~ on X such that $\pi_0(X) = X/\sim$, and prove that it is indeed an equivalence relation. (8 marks)
- (b) Let $f: X \to Y$ be a continuous map. Define the function $f_*: \pi_0(X) \to \pi_0(Y)$, and check that it is well-defined. (5 marks)
- (c) Suppose that Y is path-connected and X is not. Show that there do not exist maps $f: X \to Y$ and $g: Y \to X$ such that gf is homotopic to the identity map id_X . (6 marks)
- (d) Put $X = \{A \in M_2 \mathbb{R} \mid A^2 = A\}$. What can you say about det(A) when $A \in X$? Show that X is not path-connected. (6 marks)

(11) 2021-22 Mock Q2:

- (a) Let X be a topological space. Define the equivalence relation ~ on X such that $\pi_0(X) = X/\sim$, and prove that it is an equivalence relation. (6 marks)
- (b) Let $f: X \to Y$ be a continuous map. Define the induced map $f_*: \pi_0(X) \to \pi_0(Y)$, and prove that it is well-defined. (4 marks)
- (c) Show that if $f, g: X \to Y$ are homotopic maps then $f_* = g_*: \pi_0(X) \to \pi_0(Y)$. (4 marks)
- (d) Let Y and Z be topological spaces. Construct a bijection $\pi_0(Y \times Z) \to \pi_0(Y) \times \pi_0(Z)$, and prove that it is a bijection. (5 marks)
- (e) Define $i: \mathbb{Z} \to \mathbb{R} \setminus \mathbb{Z}$ by $i(n) = n + \frac{1}{2}$. Prove that there do not exist continuous maps $\mathbb{Z} \xrightarrow{f} S^2 \times S^2 \xrightarrow{g} \mathbb{R} \setminus \mathbb{Z}$ such that i is homotopic to $g \circ f$. (6 marks)

(12) 2023-24 Q2:

- (a) Let X be a topological space. Define the equivalence relation \sim on X such that $\pi_0(X) = X/\sim$, and prove that it is indeed an equivalence relation. (8 marks)
- (b) Let $f: X \to Y$ be a continuous map. Define the function $f_*: \pi_0(X) \to \pi_0(Y)$, and check that it is well-defined. (5 marks)
- (c) Suppose that Y is path-connected and X is not. Show that there do not exist continuous maps $f: X \to Y$ and $g: Y \to X$ such that gf is homotopic to the identity map id_X . (6 marks)
- (d) Put $X = \{A \in M_2 \mathbb{R} \mid A^2 = A\}$ (where $M_2 \mathbb{R}$ is the space of 2×2 real matrices). What can you say about det(A) when $A \in X$? Show that X is not path-connected. (6 marks)

3 The fundamental group

These questions involve material that is not covered in the current version of the course.

(13)

- (a) Let X be a topoological space, and let x_0 and x_1 be points in X. What does it mean to say that two paths from x_0 to x_1 are *pinned homotopic*? Define the set $\pi_1(X; x_0, x_1)$.
- (b) Let X be path-connected. Prove that the group $\pi_1(X; x_0)$ is isomorphic to the group $\pi_1(X; x_1)$.
- (c) Put $X = \{(w, x, y, z) \in \mathbb{C}^4 \mid w \neq x, x \neq y, y \neq z\}$, and take $x_0 = (0, 1, 2, 3)$ as the basepoint in X. Calculate $\pi_1(X)$. (You may wish to consider the expression f(w, x, y, z) = (w, x w, y x, z y).)

(14)

- (a) Let X be a based topological space, and let Y be a subspace of X containing the basepoint. What does it mean to say that Y is a *retract* of X?
- (b) Prove that if Y is a retract of X, then $|\pi_1(Y)| \le |\pi_1(X)|$.
- (c) Recall that $\mathbb{R}P^3$ is a subspace of the space $M_4(\mathbb{R})$ of all 4×4 matrices over \mathbb{R} , which is homeomorphic to \mathbb{R}^{16} . Prove that $\mathbb{R}P^3$ is not a retract of $M_4(\mathbb{R})$.
- (d) Recall that U(3) is the space of 3×3 matrices A over \mathbb{C} such that $A^{\dagger}A = I$. You may assume that for such A we have $\det(A) \in S^1$. Define $j: S^1 \to U(3)$ by

$$j(z) = \left(\begin{array}{rrr} z & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array}\right).$$

What is det(j(z))? Deduce that $\pi_1(U(3))$ is infinite.

4 Homotopy equivalence

(15)

- (a) Let $f, g: X \to Y$ be continuous maps between topological spaces. What does it mean to say that f is homotopic to g?
- (b) Let X and Y be topological spaces. What does it mean to say that X and Y are homotopy equivalent?
- (c) Show that if X and Y are homotopy equivalent then there is a bijection between the sets of path-components $\pi_0(X)$ and $\pi_0(Y)$.
- (d) Consider the cross $X = \{(x, 0) \mid -1 \le x \le 1\} \cup \{(0, y) \mid -1 \le y \le 1\}$, and let $C = \mathbb{R}^2 \setminus X$ be its complement. Prove that C is homotopy equivalent to S^1 .
- (16) Consider a metric space X.
 - (a) (i) What does it mean to say that a subset U of X is open?(ii) What does it mean to say that a subset F of X is closed?
 - (b) Show that a subset $F \subseteq X$ is closed iff for every sequence (x_n) in F that converges to a point $x \in X$, we actually have $x \in F$.
 - (c) Explain what it means for a subset $A \subseteq X$ to be compact. Show that if A is compact and $f: X \to Y$ is continuous then f(A) is compact.
 - (d) Prove that the space [0,1] is compact. Show that there is a continuous bijection $g: [-1, -1/2) \cup [1/2, 1] \rightarrow [0, 1]$; can it be chosen to be a homeomorphism?

(17)

- (a) What does it mean to say that a topological space X is homotopy equivalent to a topological space Y? Show that the relation of homotopy equivalence is an equivalence relation.
- (b) What does it mean for a space to be (a) *contractible* and (b) *path connected*? Show that any contractible space is path connected. Is the reverse implication true?
- (c) Consider the rational comb space

$$X = \{ (x, y) \in \mathbb{R}^2 \mid y \ge 0 \text{ or } x \in \mathbb{Q} \}.$$

Show that X is homotopy equivalent to the upper half plane $Y = \{(x, y) \in \mathbb{R}^2 \mid y \ge 0\}$, and deduce that X is contractible.

(18)

- (a) What does it mean to say that a topoological space X is homotopy equivalent to a topoological space Y? Show that the relation of homotopy equivalence is an equivalence relation.
- (b) What does it mean for a space to be (i) *contractible* and (ii) *path connected*? Show that any contractible space is path connected. Is the reverse implication true?
- (c) Consider the rational comb space

$$X = \{ (x, y) \in \mathbb{R}^2 \mid y \ge 0 \text{ or } x \in \mathbb{Q} \}.$$

Show that X is homotopy equivalent to the upper half plane $Y = \{(x, y) \in \mathbb{R}^2 \mid y \ge 0\}$, and deduce that X is contractible.

(19)

- (a) Let X be a subspace of \mathbb{R}^n , and let a be a point in X.
 - (i) Explain what it means for X to be *star-shaped* around a. (4 marks)
 - (ii) Prove that if X is star-shaped around a, then X is contractible. (4 marks)
- (b) (i) Suppose that $\alpha, \beta > 0$ and that $0 \le t \le 1$. Show that $\alpha t + \beta(1-t)$ is strictly greater than zero. (3 marks)
 - (ii) Suppose that $\gamma, \delta, \epsilon > 0$ and that $0 \le t \le 1$. Show that $\gamma t^2 + \delta t(1-t) + \epsilon(1-t)^2$ is strictly greater than zero. (3 marks)
 - (iii) Consider a matrix $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M_2 \mathbb{R}$. Put $\lambda = \text{trace}(A)$ and $\mu = \det(A)$. Express trace((1-t)I + tA) and $\det((1-t)I + tA)$ in terms of λ , μ and t. (6 marks)
 - (iv) Put $X = \{A \in M_2 \mathbb{R} \mid \det(A) > 0 \text{ and } \operatorname{trace}(A) > 0\}$. Prove that X is contractible. (5 marks)

(20)

- (a) Given topoological spaces X, Y and continuous maps $f, g: X \to Y$, what does it mean for f and g to be homotopic? (3 marks)
- (b) Show that if Y is contractible, then any two maps $f, g: X \to Y$ are homotopic. (7 marks)
- (c) Show that if X is contractible and Y is path-connected, then any two maps $f, g: X \to Y$ are homotopic. (10 marks)
- (d) Regard S^1 as $\{z \in \mathbb{C} \mid |z| = 1\}$, and put $T = S^1 \times S^1$. Define $f: T \to T$ by f(z, w) = (iz, -iw). Prove that f is homotopic to the identity map. (5 marks)
- (21) Let E be the figure eight space, so $E = E_{-} \cup E_{+}$ where E_{\pm} is the circle of radius one centred at $(\pm 1, 0)$.
 - (a) Prove that E is not homotopy equivalent to the torus. (4 marks)
- (b) Put $A = \{(1,0), (-1,0)\}$ and $X = \mathbb{R}^2 \setminus A$. Sketch a proof that X is homotopy equivalent to E. (5 marks)
- (c) Put $B = \{(x, y, z) \in \mathbb{R}^3 \mid x^2 = 1, y = 0\}$ and $Y = \mathbb{R}^3 \setminus B$. Deduce that Y is homotopy equivalent to E. (4 marks)
- (d) Put $C = \{(x, y, z) \in \mathbb{R}^3 \mid x^2 = 1, y = xz\}$ and $Z = \mathbb{R}^3 \setminus C$. Deduce that Z is homotopy equivalent to E. You may wish to consider the expression

$$(x, \operatorname{rot}_{\pi x/4}(y, z)) = (x, \cos(\pi x/4)y - \sin(\pi x/4)z, \sin(\pi x/4)y + \cos(\pi x/4)z).$$

(12 marks)

5 Abelian groups and chain complexes

(22)

- (a) In the context of Abelian groups, define the terms
 - homomorphism (2 marks)
 - subgroup (2 marks)
 - kernel (2 marks)
 - image. (2 marks)
- (b) Let A and B be Abelian groups, and let $\phi: A \to B$ be a homomorphism. Prove that

- (i) The kernel of ϕ is a subgroup of A (3 marks)
- (ii) The kernel of ϕ is a subgroup of the kernel of the homomorphism 2ϕ . (2 marks)
- (c) Let $\phi: \mathbb{Z} \times \mathbb{Z} \to \mathbb{Z} \times \mathbb{Z}/12$ be the homomorphism defined by

$$\phi(n,m) = (3n, (2n + 4m \mod 12)).$$

Give an isomorphism $\psi \colon \mathbb{Z} \to \ker(\phi)$. (6 marks)

(d) Let A be a finite Abelian group, and let B be a free Abelian group. Prove that if $\phi: A \to B$ is a homomorphism, then $\phi = 0$. (6 marks)

(23) 2018-19 Q4: Let $U_* \xrightarrow{i} V_* \xrightarrow{p} W_*$ be a short exact sequence of chain complexes and chain maps.

(a) Define what is meant by saying that the above sequence is short exact. (3 marks)

Now recall that a *snake* for the above sequence is a system (c, w, v, u, a) such that

- $c \in H_n(W);$
- $w \in Z_n(W)$ is a cycle such that c = [w];
- $v \in V_n$ is an element with p(v) = w;
- $u \in Z_{n-1}(U)$ is a cycle with $i(u) = d(v) \in V_{n-1}$;
- $a = [u] \in H_{n-1}(U).$
- (b) Prove that for each $c \in H_n(W)$ there is a snake starting with c. (8 marks)
- (c) Prove that if two snakes have the same starting point, then they also have the same endpoint. (10 marks)
- (d) Suppose that the differential $d: V_{n+1} \to V_n$ is surjective. Show that any snake starting in $H_n(W)$ ends with zero. (4 marks)

(24) 2021-22 Mock Q3:

- (a) Let $U_* \xrightarrow{i} V_* \xrightarrow{p} W_*$ be a short exact sequence of chain complexes and chain maps. Define what is meant by a *snake* for this sequence. (5 marks)
- (b) Define the homomorphism $\delta: H_n(W) \to H_{n-1}(U)$. You should give a clear statement of the lemmas needed to ensure that your definition is meaningful, but you do not need to prove those lemmas. (4 marks)
- (c) Suppose that $H_k(W)$ is finite for all k, and that $H_k(U) \simeq \mathbb{Z}$ for all k. Prove that $H_k(V)$ is infinite and that the map $p_*: H_k(V) \to H_k(W)$ is surjective. (5 marks)
- (d) Consider the chain complex with $A_k = \mathbb{Z}^3$ for all $k \in \mathbb{Z}$ and d(x, y, z) = (z, 0, 0).
 - (i) Find the homology of A_* . (2 marks)
 - (ii) Show that the formula m(x, y, z) = (0, y, 0) defines a chain map $m: A_* \to A_*$ (2 marks)
 - (iii) Show that m is chain homotopic to the identity. (3 marks)
 - (iv) Construct a chain complex A'_* where the differential is zero, and a chain homotopy equivalence from A'_* to A_* . (4 marks)

(25) 2021-22 Q3:

(a) Define the terms chain complex, chain map and chain homotopy. (8 marks)

- (b) Prove that if two chain maps are chain homotopic, then they have the same effect on homology groups. (5 marks)
- (c) Consider the chain complex T with T_i = Z/8 for all i and d(x) = 4x for all x. Find the homology groups of T.
 (3 marks)
- (d) Suppose we have a short exact sequence $A_* \xrightarrow{i} B_* \xrightarrow{p} C_*$ of chain complexes and chain maps. Suppose that for all $i \in \mathbb{Z}$ we have $H_{2i+1}(A) = H_{2i+1}(C) = 0$ and $|H_{2i}(A)| = 3$ and $|H_{2i}(C)| = 5$. Prove that all homology groups of B are cyclic or trivial, and determine their orders. (5 marks)
- (e) Let U_* be a chain complex in which all the differentials d_{2i} (for all $i \in \mathbb{Z}$) are surjective homomorphisms. What can we conclude about the homology groups of U_* ? (4 marks)

(26) 2022-23 Q3: Let $U_* \xrightarrow{i} V_* \xrightarrow{p} W_*$ be a short exact sequence of chain complexes and chain maps.

(a) Define what is meant by saying that the above sequence is short exact. (3 marks)

Now recall that a *snake* for the above sequence is a system (c, w, v, u, a) such that

- $c \in H_n(W);$
- $w \in Z_n(W)$ is a cycle such that c = [w];
- $v \in V_n$ is an element with p(v) = w;
- $u \in Z_{n-1}(U)$ is a cycle with $i(u) = d(v) \in V_{n-1}$;

•
$$a = [u] \in H_{n-1}(U).$$

- (b) Prove that for each $c \in H_n(W)$ there is a snake starting with c. (7 marks)
- (c) Explain how the connecting homomorphism $\delta: H_n(W) \to H_{n-1}(U)$ is defined in terms of snakes. If any further lemmas are needed to ensure that your definition is meaningful, then you should state those lemmas carefully, but you need not prove them. (4 marks)
- (d) Consider the following example. For each $k \in \mathbb{Z}$ we have

$U_k = \mathbb{Z}/24 = \mathbb{Z}/(2^3 \times 3)$	$d^U(x) = 12x = 2^2 \times 3 \times x$
$V_k = \mathbb{Z}/1296 = \mathbb{Z}/(2^4 \times 3^4)$	$d^V(x) = 36x = 2^2 \times 3^2 \times x$
$W_k = \mathbb{Z}/54 = \mathbb{Z}/(2 \times 3^3)$	$d^W(x) = -18x = -2 \times 3^2 \times x.$

The maps

$$U_k \xrightarrow{i} V_k \xrightarrow{p} W_k$$

are $i(a \pmod{24}) = 54a \pmod{1296}$ and $p(b \pmod{1296}) = b \pmod{54}$.

- (i) Check that i and p are chain maps. (You may assume that they give a short exact sequence.) (3 marks)
- (ii) Calculate the groups $H_k(U)$, $H_k(V)$ and $H_k(W)$. (5 marks)
- (iii) By finding an appropriate snake, calculate the homomorphism $\delta: H_k(W) \to H_{k-1}(U)$. (3 marks)

(27)

- (a) Define the terms chain map, chain homotopy, chain homotopic and chain homotopy equivalence. (8 marks)
- (b) Show that if $f, g: U_* \to V_*$ are chain maps that are chain homotopic to each other, then $f_* = g_*: H_*(U) \to H_*(V)$. (5 marks)
- (c) Consider the chain complex T_* with $T_i = \mathbb{Z}^2$ for all i and $d_i(x, y) = (y, 0)$ for all $(x, y) \in T_i$. Show that T_* is chain homotopy equivalent to the zero complex. (4 marks)

- (d) Suppose we have a short exact sequence $A_* \xrightarrow{i} B_* \xrightarrow{p} C_*$ of chain complexes and chain maps. Suppose that for all $k \in \mathbb{Z}$ we have $H_k(B) = 0$. Suppose also that $H_k(A) = \mathbb{Z}/2^k$ for $k \ge 0$ and $H_k(A) = 0$ for k < 0. Determine the homology groups of C_* . (3 marks)
- (e) Let U_* be a chain complex in which $U_k = 0$ for k < 0 and $|U_k| = 2^k$ for $k \ge 0$ and $d_{2i}: U_{2i} \to U_{2i-1}$ is surjective for all *i*. Find the homology groups of U_* . (5 marks)

(28) 2023-24 Q3:

- (a) Define the terms chain map, chain homotopy, chain homotopic and chain homotopy equivalence. (8 marks)
- (b) Show that if $f, g: U_* \to V_*$ are chain maps that are chain homotopic to each other, then $f_* = g_*: H_*(U) \to H_*(V)$. (5 marks)
- (c) Consider the chain complex T_* with $T_i = \mathbb{Z}^2$ for all i and $d_i(x, y) = (y, 0)$ for all $(x, y) \in T_i$. Show that T_* is chain homotopy equivalent to the zero complex. (4 marks)
- (d) Suppose we have a short exact sequence $A_* \xrightarrow{i} B_* \xrightarrow{p} C_*$ of chain complexes and chain maps. Suppose that for all $k \in \mathbb{Z}$ we have $H_k(B) = 0$. Suppose also that $H_k(A) = \mathbb{Z}/2^k$ for $k \ge 0$ and $H_k(A) = 0$ for k < 0. Determine the homology groups of C_* . (3 marks)
- (e) Let U_* be a chain complex in which $U_k = 0$ for k < 0 and $|U_k| = 2^k$ for $k \ge 0$ and $d_{2i}: U_{2i} \to U_{2i-1}$ is surjective for all *i*. Find the homology groups of U_* . (5 marks)

6 Singular chains

- (29) Let X be a topological space.
 - (a) Let $c: \Delta_1 \to X$ be a constant path. Prove that c is homologous to 0.
 - (b) Let $s: \Delta_1 \to X$ be a path. Define the reversed path \overline{s} , and prove that \overline{s} is homologous to -s.
 - (c) Let $r, s: \Delta_1 \to X$ be paths such that $r(e_1) = s(e_0)$. Write down a path $u: \Delta_1 \to X$ and prove that u is homologous to r + s.
 - (d) Let X be the complement of the shaded disc in the diagram below. Write down a path $u: \Delta_1 \to X$ such that u is homologous to 2p 2q 2r + s.



(30)

(a) Let X be a topological space.

- (i) Define the groups $C_n(X)$ for all nonnegative integers n. (2 marks)
- (ii) Define the homomorphisms ∂_n . (3 marks)
- (iii) Prove that $\partial_1 \circ \partial_2 = 0$. (3 marks)
- (iv) Define the groups $H_n(X)$. (4 marks)
- (b) Describe (without proof, but with careful attention to any special cases) the groups $H_n(\mathbb{R}^k \setminus \{0\})$ for all $n \ge 0$ and all $k \ge 1$. (5 marks)
- (c) Let $u = n_1 s_1 + \ldots + n_k s_k$ be an *m*-cycle in S^n (where m > 0), and suppose that there is a point $a \in S^n$ that is not contained in any of the sets $s_1(\Delta_m), \ldots, s_k(\Delta_m)$. Prove that u is a boundary. (8 marks)

(31)

- (a) Let X be a topological space.
 - (i) Define the groups $C_0(X)$ and $C_1(X)$, and the homomorphism $\partial_1 : C_1(X) \to C_0(X)$.
 - (ii) Define the subdivision homomorphism sd: $C_1(X) \to C_1(X)$.
 - (iii) Prove that $\partial_1 \operatorname{sd}^n(u) = \partial_1(u)$ for all $n \ge 1$.
 - (iv) Prove that if $u \in B_1(X)$ then $sd(u) \in B_1(X)$.
 - (v) Let A and B be points in a vector space V. Give an expression for $sd\langle A, B \rangle$ in terms of paths of the form $\langle C, D \rangle$.
- (b) Describe without proof the groups $H_1(S^1)$, $H_1(S^1 \times S^1)$, $H_1(\mathbb{R}P^2)$ and $H_1(\mathbb{R}^3 \setminus \{0\})$.
- (c) For each element $u \in H_1(\mathbb{R}P^2)$, give a path s in $\mathbb{R}P^2$ such that u = [s].
- (32) Consider the following diagram.



Let X be the complement in \mathbb{R}^2 of the shaded disc. Define $u, v, w \in C_1(X)$ by

$$\begin{split} & u = \langle A, B \rangle + \langle B, E \rangle + \langle E, A \rangle \\ & v = \langle A, B \rangle + \langle B, C \rangle + \langle C, D \rangle + \langle D, A \rangle \\ & w = \langle A, E \rangle + \langle E, D \rangle + \langle D, A \rangle. \end{split}$$

- (a) Prove that u is a cycle. (2 marks)
- (b) Prove that $\langle B, B \rangle$ is homologous to 0 in X. (3 marks)
- (c) Prove that $\langle E, B \rangle$ is homologous to $-\langle B, E \rangle$ in X. (4 marks)
- (d) Prove in detail that u is homologous to v in X, justifying each step. (8 marks)
- (e) Write down a basic 1-chain s that is homologous in X to $\langle A, B \rangle + \langle B, C \rangle$ (5 marks)
- (f) Is *u* homologous to *w*? Give a brief reason for your answer. (3 marks)

(33) 2023-24 Q4:

- (a) Let X be a topological space.
 - (i) Define the groups $C_n(X)$ for all nonnegative integers n. (2 marks)
 - (ii) Define the homomorphisms $\partial_n : C_n(X) \to C_{n-1}(X)$. (3 marks)
 - (iii) Prove that $\partial_1 \circ \partial_2 = 0$. (3 marks)
 - (iv) Define the groups $H_n(X)$. (4 marks)
- (b) Describe (without proof, but with careful attention to any special cases) the groups $H_n(\mathbb{R}^k \setminus \{0\})$ for all $n \ge 0$ and all $k \ge 1$. (5 marks)
- (c) Let $u = n_1 s_1 + \ldots + n_k s_k$ be an element of $Z_m(S^n)$ (where m > 0), and suppose that there is a point $a \in S^n$ that is not contained in any of the sets $s_1(\Delta_m), \ldots, s_k(\Delta_m)$. Prove that u is a boundary. (You may assume standard results and calculations from the course so long as you state them carefully.) (8 marks)

7 True or false

(34) Are the following statements true or false? Give proof or disproof as appropriate. You may quote general theorems, provided that you state them clearly.

- (a) The punctured disc $X = \{(x, y) \in \mathbb{R}^2 \mid 0 < x^2 + y^2 \le 1\}$ is compact.
- (b) The circle S^1 is homeomorphic to $S^1 \times I$.
- (c) The circle S^1 is homotopy equivalent to $S^1 \times I$.
- (d) $\mathbb{C} \setminus S^1$ is homotopy equivalent to $Y = \{z \in \mathbb{C} \mid z = 0 \text{ or } |z| = 1\}.$
- (e) Every continuous bijection from $[0,1] \cup (2,3]$ to [0,1] is a homeomorphism.

(35) Are the following statements true or false? Give proof or disproof as appropriate. You may quote general theorems, provided that you state them clearly.

- (a) S^3 is contractible.
- (b) If a space X is the union of two closed, path-connected subspaces A and B, then X is path-connected.
- (c) $(\mathbb{R} \times \mathbb{R}) \setminus (\mathbb{R} \times \{0\})$ is homotopy equivalent to S^1 .
- (d) $(\mathbb{R} \times \mathbb{R}^2) \setminus (\mathbb{R} \times \{0\})$ is homotopy equivalent to S^1 .
- (e) The space $\mathbb{C} \setminus \{0, 1\}$ is homeomorphic to $\mathbb{C} \setminus \{i, -i\}$.
- (f) The space $\mathbb{C} \setminus \{0, 1\}$ is homotopy equivalent to $\mathbb{C} \setminus \{0, 1, 2\}$.

(36) Are the following statements true or false? Give proof or disproof as appropriate. You may quote general theorems, provided that you state them clearly.

- (a) The identity map of the unit circle is homotopic to the constant map $c: S^1 \to S^1$ defined by c(z) = 1 for all z.
- (b) Let $f_n: S^1 \to S^1$ be defined by $f_n(z) = z^n$. Then f_n is not homotopic to f_m when $n \neq m$.
- (c) \mathbb{R}^2 is homeomorphic to \mathbb{R}^3 .
- (d) If $f: X \to X$ is a homotopy equivalence, then $f_*: H_1(X) \to H_1(X)$ is the identity map.

(37) Are the following statements true or false? Give proof or disproof as appropriate. You may quote general theorems, provided that you state them clearly.

- (a) The torus $T = S^1 \times S^1$ is homotopy equivalent to S^2 .
- (b) There is a map $r: B^4 \to S^3$ such that rj is homotopic to id_{S^3} , where $j: S^3 \to B^4$ is the inclusion map.
- (c) \mathbb{R}^2 is homeomorphic to \mathbb{R}^3 .
- (d) Every continuous function $f: S^2 \to \mathbb{R}^3$ is homotopic to a constant function.
- (e) Let $K \subset S^3$ be a trefoil knot. Then $S^3 \setminus K$ is homotopy equivalent to $\mathbb{R}P^2$.

(38) Are the following statements true or false? Give proof or disproof as appropriate. You may quote general theorems or calculations of homology groups without proof, provided that you state them clearly. You may give pictures instead of formulae, provided that they are clearly explained.

- (a) $\mathbb{R}P^1$ is homeomorphic to S^1 .
- (b) The Möbius strip is homotopy equivalent to S^2 .
- (c) $S^2 \setminus S^1$ is homotopy equivalent to $\mathbb{R} \setminus \{0\}$.
- (d) The letter A is homeomorphic to the letter D.
- (e) Any compact convex subset of \mathbb{R}^2 is homeomorphic to B^2 .

(39) Are the following statements true or false? Give proof or disproof as appropriate. You may quote general theorems or calculations of homology groups without proof, provided that you state them clearly. You may give pictures instead of formulae, provided that they are clearly explained.

- (a) $\mathbb{R}P^1$ is homeomorphic to S^1 .
- (b) The Möbius strip is homotopy equivalent to S^2 .
- (c) SO(3) is homeomorphic to $\mathbb{R}P^3$.
- (d) $S^2 \setminus S^1$ is homotopy equivalent to $\mathbb{R} \setminus \{0\}$.
- (e) The letter A is homeomorphic to the letter D.
- (f) Any compact convex subset of \mathbb{R}^2 is homeomorphic to B^2 .

(40) Are the following statements true or false? Give proof or disproof as appropriate. You may quote general theorems or calculations of homology groups without proof, provided that you state them clearly. You may give pictures instead of formulae, provided that they are clearly explained.

- (a) S^1 is homotopy equivalent to S^2 . (3 marks)
- (b) S^1 is homotopy equivalent to the Möbius strip. (4 marks)
- (c) S^1 is homeomorphic to the Möbius strip. (4 marks)
- (d) $\mathbb{R}P^2$ is homeomorphic to $S^1 \times S^1$. (4 marks)
- (e) $SU(2) \setminus \{I\}$ is homeomorphic to \mathbb{R}^3 . (5 marks)
- (f) $\Delta_n \times \Delta_m$ is homeomorphic to Δ_{n+m} . (5 marks)

(41) Are the following statements true or false? Give proof or disproof as appropriate. You may quote general theorems without proof, provided that you state them clearly. You may give pictures instead of formulae, provided that they are clearly explained.

- (a) There is a continuous surjective map from $S^1 \times S^1$ to \mathbb{R}
- (b) $\mathbb{C} \setminus \{2\}$ is homotopy equivalent to S^1
- (c) $\mathbb{C} \setminus \{-1, 1\}$ is homotopy equivalent to S^1
- (d) $S^2 \setminus \{\text{the north pole}\}\$ is homeomorphic to \mathbb{C} .
- (e) The letter X (considered as a subspace of \mathbb{R}^2) is homeomorphic to the letter Y.
- (f) The letter X (considered as a subspace of \mathbb{R}^2) is homotopy equivalent to the letter Y.

(42) Are the following statements true or false? Give proof or disproof as appropriate. You may quote general theorems without proof, provided that you state them clearly. You may give pictures instead of formulae, provided that they are clearly explained.

- (a) There is a continuous surjective map from $\mathbb{R} \times \mathbb{R}$ to $\mathbb{R} \setminus \{0\}$ (5 marks)
- (b) $S^2 \setminus S^1$ is homeomorphic to \mathbb{R}^2 (5 marks)
- (c) SO(2) is homotopy equivalent to the Möbius strip (5 marks)
- (d) SO(3) is homotopy equivalent to the torus (5 marks)
- (e) The space $X = S^1 \cup \{(x,0) \mid x \in \mathbb{R}\}$ is homeomorphic to $Y = S^1 \cup \{(x,1) \mid x \in \mathbb{R}\}$. (5 marks)

(43) Are the following statements true or false? Give proof or disproof as appropriate. You may quote general theorems without proof, provided that you state them clearly. You may give pictures instead of formulae, provided that they are clearly explained.

- (a) If X and Y are both path-connected subsets of \mathbb{R}^2 , then $X \cap Y$ is also path-connected. (5 marks)
- (b) The torus is homotopy equivalent to S^2 . (5 marks)
- (c) If $f: X \to Y$ and $g: Y \to X$ are continuous, based maps and $gf = id_X$ then $\pi_1(X) \simeq \pi_1(Y)$. (5 marks)
- (d) If two letters of the alphabet, considered as subspaces of \mathbb{R}^2 , both have infinite H_1 , then they are homotopy equivalent. (5 marks)
- (e) The space $GL_3(\mathbb{R})$ is path-connected. (5 marks)

(44) 2018-19 Q2: Are the following true or false? Justify your answers.

- (a) S^5 is a Hausdorff space. (4 marks)
- (b) The Klein bottle is a retract of $S^1 \times S^1 \times S^1$. (4 marks)
- (c) There is a connected space X with $\pi_1(X) \simeq \mathbb{Z}/2$ and $H_1(X) \simeq \mathbb{Z}$. (4 marks)
- (d) There is a short exact sequence $\mathbb{Z}/9 \to \mathbb{Z}/99 \to \mathbb{Z}/11$. (4 marks)
- (e) If K is a simplicial complex and L is a subcomplex and $H_3(K) = 0$ then $H_3(L) = 0$. (4 marks)
- (f) If K and L are simplicial complexes and $f: |K| \to |L|$ is a continuous map then there is a simplicial map $s: K \to L$ such that f is homotopic to |s|. (5 marks)

(45) 2023-24 Q5: Are the following statements true or false? Give proof or disproof as appropriate. You may quote general theorems and calculations, provided that you state them clearly.

(a) S^3 is contractible. (3 marks)

- (b) $\mathbb{R}P^3$ is a homotopy retract of S^3 . (3 marks)
- (c) If a space X is the union of two closed, path-connected subspaces A and B, then X is path-connected. (3 marks)
- (d) $(\mathbb{R} \times \mathbb{R}) \setminus (\mathbb{R} \times \{0\})$ is homotopy equivalent to S^1 . (4 marks)
- (e) $(\mathbb{R} \times \mathbb{R}^2) \setminus (\mathbb{R} \times \{0\})$ is homotopy equivalent to S^1 . (4 marks)
- (f) The space $\mathbb{C} \setminus \{0, 1\}$ is homeomorphic to $\mathbb{C} \setminus \{i, -i\}$. (4 marks)
- (g) The space $\mathbb{C} \setminus \{0, 1\}$ is homotopy equivalent to $\mathbb{C} \setminus \{0, 1, 2\}$. (4 marks)

8 Examples

- (46) Give examples of the following things, with careful justification.
- (a) A noncompact topoological space X with a sequence of compact subspaces $Y_1 \subset Y_2 \subset \ldots$ such that the union of all the sets Y_n is equal to X.
- (b) A topoological space X with two noncompact subsets Y, Z such that $Y \cup Z$ is compact.
- (c) A sequence in \mathbb{R} with no convergent subsequence.
- (d) A non-surjective map $f: X \to Y$ such that $f_*: \pi_0(X) \to \pi_0(Y)$ is surjective.
- (e) An injective map $f: X \to Y$ such that $f_*: \pi_0(X) \to \pi_0(Y)$ is not injective.
- (47) Give examples of the following things, with careful justification.
 - (a) A continuous bijection that is not a homeomorphism. (3 marks)
 - (b) An infinite sequence of open sets whose intersection is not open. (3 marks)
 - (c) Two metric spaces X, Y such that X is bounded, Y is unbounded, and X is homeomorphic to Y. (4 marks)
 - (d) A sequence in (0,1) such that no subsequence converges in (0,1). (5 marks)
 - (e) Two contractible subsets of \mathbb{R}^2 whose intersection is not contractible. (5 marks)
 - (f) Two topoological spaces X, Y and points $x \in X, y \in Y$ such that X is homotopy equivalent to Y but $X \setminus \{x\}$ is not homotopy equivalent to $Y \setminus \{y\}$. (5 marks)
- (48) Give examples of the following things, with justification.
 - (a) Connected sets $X, Y \subseteq \mathbb{R}^2$ such that $X \cap Y$ is not connected.
 - (b) A sequence of open sets $U_n \subseteq \mathbb{R}$ such that the set $X = U_1 \cap U_2 \cap \ldots = \bigcap_n U_n$ is not open.
 - (c) A surjective map $f: X \to Y$ of topological spaces such that the homomorphism $f_*: H_1(X) \to H_1(Y)$ is not surjective.
 - (d) A path connected space X that is homotopy equivalent to $X \times X$.
 - (e) A path connected space X that is not homotopy equivalent to $X \times X$.
- (49) Give examples of the following things.
- (a) A space X and a point $x \in X$ such that X is not contractible but $X \setminus \{x\}$ is contractible. (3 marks)

- (b) A subspace $X \subseteq \mathbb{R}^2$ that is homotopy equivalent to $S^4 \setminus S^2$. (You need not give a proof.) (4 marks)
- (c) Spaces X and Y, a discontinuous map $f: X \to Y$, and an open subset $V \subseteq Y$ such that $f^{-1}V$ is not open in X. (You should justify your answer carefully.) (6 marks)
- (d) A space X and a point $x \in X$ such that $\pi_1(X)$ is abelian and $\pi_1(X \setminus \{x\})$ is nonabelian. (You should state what $\pi_1(X)$ and $\pi_1(X \setminus \{x\})$ are, but no further justification is required.) (6 marks)
- (e) A space X such that a(X) = 2 and b(X) = 2, where as usual

 $a(X) = \max\{|S| \mid S \text{ is a finite subset of } X \text{ and } X \setminus S \text{ is path-connected } \}$ = the largest number of points that can be removed from X without disconnecting it

- $b(X) = \min\{|S| \mid S \text{ is a finite subset of } X \text{ and } X \setminus S \text{ is not path-connected } \}$
 - = the smallest number of points that have to be removed from X to disconnect it

(You should justify your answer, but complete rigour is not required.) (6 marks)

- (50) Give one example of each of the following things, with justification.
 - (a) A path connected space X with $H_1(X) = \mathbb{Z} \oplus (\mathbb{Z}/2)$. (4 marks)
 - (b) A path-connected space X and points $a, b, c \in X$ such that $X \setminus \{a, b, c\}$ is still path-connected. (3 marks)
 - (c) A path-connected space X and a point $a \in X$ such that $H_1(X)$ and $H_1(X \setminus \{a\})$ are both trivial. (5 marks)
 - (d) A continuous, surjective map $f: X \to Y$, where Y is compact but X is not. (3 marks)
 - (e) A space X and points $a, b \in X$ such that $\pi_1(X)$ is nonabelian but the space $Y = X \setminus \{a, b\}$ is simply connected. (5 marks)
 - (f) A continuous bijection that is not a homeomorphism. (5 marks)

(51) 2020-21 Q1: Give examples as follows, justifying your answers.

- (a) Topological spaces X and Y, together with injective functions $f: X \to Y$ and $g: Y \to X$ such that $f, f \circ g$ and $g \circ f$ are all continuous, but g is not continuous. (4 marks)
- (b) A compact, path-connected space X together with a continuous map $f: X \to X$ with no fixed points. (4 marks)
- (c) A space X such that $H_1(X)$ is not a free abelian group. (Note here that the zero group is free abelian with no generators, so in particular $H_1(X)$ must be nonzero.) (4 marks)
- (d) A space X together with points $a, b, c \in X$ such that $|\Pi(X; a, b)| \neq |\Pi(X; b, c)|$. (4 marks)
- (e) A space X such that $\pi_1(X)$ is a free group with 3 generators, and $H_2(X) = \mathbb{Z}$. (4 marks)

(52) 2021-22 Mock Q4: For each of the following, either give an example (with justification) or prove that no example can exist.

- (a) A continuous map $f: X \to Y$ such that $f_*: H_1(X) \to H_1(Y)$ is injective but not surjective, and $f_*: H_{10}(X) \to H_{10}(Y)$ is surjective but not injective. (5 marks)
- (b) A path connected space X that is homotopy equivalent to $X \times X$. (5 marks)
- (c) A path connected space X that is not homotopy equivalent to $X \times X$. (5 marks)
- (d) A space X and a point $x \in X$ such that X is not contractible but $X \setminus \{x\}$ is contractible. (5 marks)

(e) A subspace $X \subseteq \mathbb{R}^2$ that is homotopy equivalent to $S^4 \setminus S^2$. (5 marks)

(53) 2021-22 Q4: For each of the following, either give an example (with justification) or prove that no example can exist.

- (a) A continuous injective map $i: X \to Y$ such that the map $i_*: H_2(X) \to H_2(Y)$ is not injective. (5 marks)
- (b) A continuous surjective map $p: X \to Y$ such that the map $p_*: H_2(X) \to H_2(Y)$ is not surjective. (5 marks)
- (c) A contractible space X and a homeomorphism $f: X \to X$ with no fixed points. (5 marks)
- (d) A continuous injective map $f: S^1 \to S^3$ such that $S^3 \setminus f(S^1)$ is homotopy equivalent to S^1 . (5 marks)
- (e) A continuous injective map $f: S^1 \to S^3$ such that $S^3 \setminus f(S^1)$ is contractible. (5 marks)

(54) 2022-23 Q4: For each of the following, either give an example (with justification) or prove that no example can exist.

- (a) A topological space X with two noncompact subsets Y, Z such that $Y \cup Z$ is compact. (5 marks)
- (b) Subsets $A, B, C \subseteq \mathbb{R}^2$ such that $A, B, C, A \cup B, A \cup C$ and $B \cup C$ are all contractible, but $A \cup B \cup C$ is not contractible. (5 marks)
- (c) A topological space X with two open subsets U and V such that U, V and $U \cap V$ are all homotopy equivalent to S^1 , and $X = U \cup V$, and X is homotopy equivalent to S^4 . (5 marks)
- (d) A path connected space X such that $H_*(X)$ is not isomorphic to $H_*(X \times X)$. (5 marks)
- (e) Spaces X and Y such that X is path connected, Y is not path connected, and $H_k(X) \simeq H_k(Y)$ for all k. (5 marks)

9 Real projective space

In the current version of the course, in the Introduction we define $\mathbb{R}P^n = S^n/(x \sim -x)$ and

$$P_n = \{A \in M_{n+1}(\mathbb{R}) \mid A^2 = A^T = A, \text{ trace}(A) = 1\},\$$

and we mention that $\mathbb{R}P^n$ is homeomorphic to P_n . A proof is given in Problem Sheet 5. In some earlier versions of the course, $\mathbb{R}P^n$ was just defined to be the same as P_n . Problems in this section should be approached from that point of view.

(55)

- (a) Define the set $\mathbb{R}P^2$ and the map $q: S^2 \to \mathbb{R}P^2$.
- (b) Define the usual metric on $\mathbb{R}P^2$, and prove that it is a metric.
- (c) Define the space Δ_2 , and prove carefully that there is a surjective continuous map $f \colon \mathbb{R}P^2 \to \Delta_2$ satisfying $fq(u, v, w) = (u^2, u^2 + v^2)$ for all $(u, v, w) \in \Delta_2$. You may use general theorems provided that you state them precisely.

(56)

- (a) Define the set $\mathbb{R}P^n$, and write down a metric on it, proving that your formula is well-defined. (You need not show that it is a metric.) (6 marks)
- (b) Define what it means for a metric space X to be sequentially compact. (3 marks)
- (c) Define the set $\pi_0(X)$, and say what it means for X to be *path-connected*. (6 marks)
- (d) Prove that the space $\mathbb{R}P^n$ is sequentially compact and path-connected. State clearly any general theorems or results that you use. (10 marks)

10 Multipart questions

(57)

- (a) What is a *metric space*? What is a *continuous function*?
- (b) Define the discrete metric on a set X.
- (c) Let X be a space with a discrete metric. Show that any path $s: \Delta_1 \to X$ is constant, and deduce that $\pi_0(X) = X$.
- (d) Consider the space $Y = \{(x, y) \in \mathbb{R}^2 \mid xy \neq 0\}$ and show that $\pi_0(Y)$ has precisely four elements. If $f: Y \to Y$ denotes reflection in the line x = y, describe the map $f_*: \pi_0(Y) \to \pi_0(Y)$. Is f homotopic to the identity map?

(58) 2018-19 Q1:

- (a) Given a topological space X, define the set $\pi_0(X)$. You should include a proof that the relevant equivalence relation is in fact an equivalence relation. (8 marks)
- (b) Consider [0, 1] as a based space with 0 as the basepoint. For $n \ge 3$ we define $X_n = \{z \in \mathbb{C} \mid z^n \in [0, 1]\}$:



- (i) For which n and m (with $n, m \ge 3$) is X_n homotopy equivalent to X_m ? (3 marks)
- (ii) For which n and m (with $n, m \ge 3$) is X_n homeomorphic to X_m ? (4 marks)

Justify your answers carefully.

- (c) Give examples as follows, with justification:
 - (1) A based space W with $|\pi_1(W)| = 8$. (3 marks)
 - (2) A space X with two points $a, b \in X$ such that $\pi_1(X, a)$ is not isomorphic to $\pi_1(X, b)$. (3 marks)
 - (3) A space Y such that H₀(Y) ≃ H₂(Y) ≃ H₄(Y) ≃ H₆(Y) ≃ Z and all other homology groups are trivial. (4 marks)

(59) 2021-22 Mock Q1: For $n \ge 3$, we put $X_n = \mathbb{R}^2 \setminus \{(1,0), (2,0), \dots, (n,0)\}.$

- (a) Define the following terms: topology, topological space, continuous map, homeomorphism. (7 marks)
- (b) Find a space Y_n consisting of a finite number of straight line segments that is homotopy equivalent to X_n . Give a brief justification for the claim that Y_n is homotopy equivalent to X_n . (6 marks)
- (c) Prove that X_n is not homeomorphic to Y_n . (3 marks)
- (d) Prove that X_n is not homotopy equivalent to S^m for any m. (4 marks)
- (e) Find contractible open sets $U_n, V_n \subseteq \mathbb{C}$ such that $X_n = U_n \cup V_n$. Give a careful proof that U_n and V_n are contractible. (5 marks)

Claims about the homology of particular spaces should be stated clearly and justified briefly, but details are not required.

(60) 2021-22 Q1: For $n \ge 3$, we put

$$X_n = \{ z \in \mathbb{C} \mid |z| = 1 \text{ or } z^n \in (0, \infty) \}$$

$$Y_n = \{ z \in \mathbb{C} \mid |z| = 1 \text{ or } z^n \in [0, \infty) \}.$$

- (a) Sketch X_3 and Y_3 . (2 marks)
- (b) Define the terms homotopy and homotopy equivalent. (5 marks)
- (c) Prove (by constructing explicit maps and homotopies, and checking their validity) that X_n and X_m are homotopy equivalent for all $n, m \ge 3$. (8 marks)
- (d) Prove that for all $n \neq m$, the space X_n is not homeomorphic to X_m . (6 marks)
- (e) Prove that for all $n \neq m$, the space Y_n is not homotopy equivalent to Y_m . (4 marks)

Claims about the homology of particular spaces should be stated clearly and justified briefly, but details are not required.

- (61) 2018-19 Q3: Let K and L be abstract simplicial complexes.
 - (a) Define what is meant by a *simplicial map* from K to L. (3 marks)
 - (b) Let $s, t: K \to L$ be simplicial maps. Define what it means for s and t to be directly contiguous. (3 marks)
 - (c) Prove that if s and t are directly contiguous, then the resulting maps $|s|, |t|: |K| \to |L|$ are homotopic. (3 marks)
 - (d) Prove that if s and t are directly contiguous, then the resulting maps $s_*, t_* \colon H_*(K) \to H_*(L)$ are the same. (You can prove the main formula just for n = 3 rather than general n.) (9 marks)
 - (e) How many injective simplicial maps are there from $\partial \Delta^2$ to itself? Show that no two of them are directly contiguous. (7 marks)

(62) 2018-19 Q5: Consider a simplicial complex K with subcomplexes L and M such that $K = L \cup M$. Use the following notation for the inclusion maps:

$$\begin{array}{ccc} L \cap M & \stackrel{i}{\longrightarrow} & L \\ j & & \downarrow^{f} \\ M & \stackrel{a}{\longrightarrow} & K. \end{array}$$

- (a) State the Seifert-van Kampen Theorem (in a form applicable to simplicial complexes and subcomplexes as above).
 (4 marks)
- (b) State the Mayer-Vietoris Theorem. (5 marks)
- (c) State a theorem about the relationship between π_1 and H_1 . (3 marks)
- (d) Suppose that |L|, |M| and $|L \cap M|$ are all homotopy equivalent to S^1 . Suppose that the maps i and j both have degree two.
 - (1) Find a presentation for $\pi_1(|K|)$. (3 marks)
 - (2) Find H_{*}(K). In particular, you should express each nonzero group as a direct sum of terms like Z or Z/n.
 (10 marks)
- (63) 2019-20 Q1: Consider the following spaces:



$$X_{6} = (S^{1} \times S^{1}) \setminus \{(1,1)\}$$

$$X_{7} = GL_{2}(\mathbb{R}) = \{A \in M_{2}(\mathbb{R}) \mid \det(A) \neq 0\}$$

$$X_{8} = \mathbb{R}$$

$$X_{9} = \{(u,v) \in \mathbb{C}^{2} \mid 1 \leq |u| \leq 2 \leq |v| \leq 3\}.$$

(Here X_3 and X_4 are closed orientable surfaces, and X_5 is the union of X_4 with a line segment with one endpoint lying on X_4 . Everything else should be clear.)

- (a) These 10 spaces can be grouped into 5 pairs $\{X_i, X_j\}$ such that X_i is homotopy equivalent to X_j . Find these pairs, and justify your answers. In each case you should prove that X_i is homotopy equivalent to X_j , and also that it is not homotopy equivalent to any of the other spaces. (25 marks)
- (b) For each pair $\{X_i, X_j\}$ as in (a), prove that X_i is not homeomorphic to X_j . (In one case you may need to appeal to some geometric intuition, but you should be able to give a more formal proof in the other four cases.) (15 marks)

(64) 2019-20 Q2:

- (a) Let A and B be finite abelian groups such that |A| and |B| are coprime.
 - (i) What can you say about homomorphisms from A to B? (10 marks)
 - (ii) Now suppose we have a short exact sequence $A \to U \to B$ of abelian groups. By considering the classification of finite abelian groups, or otherwise, what can you say about U? (15 marks)
- (b) Let X be a topological space, with open subspaces U and V such that $X = U \cup V$. Suppose that U, V, X and $U \cap V$ are all path-connected, and that for all k > 0 we have $H_k(U \cap V) = \mathbb{Z}/2^k$ and $H_k(U) = \mathbb{Z}/3^k$ and $H_k(V) = \mathbb{Z}/5^k$. Calculate $H_*(X)$. (15 marks)

(65) 2020-21 Q2: Fix $n \ge 2$. Define an equivalence relation on the disc $B^2 = \{z \in \mathbb{C} \mid |z| \le 1\}$ by $z_0 \sim z_1$ iff $z_0 = z_1$, or $(|z_0| = |z_1| = 1 \text{ and } z_0^n = z_1^n)$. Put $X = B^2 / \sim$ and

$$Y = \{(u, v) \in \mathbb{C}^2 \mid |u| \le 1, \quad v^n = (1 - |u|)^n u\}.$$

Note that when n = 2 we just have $X = \mathbb{R}P^2$; this should guide your thinking about the general case.

(a) Show carefully that there is a homeomorphism $f: X \to Y$ such that $f([z]) = (z^n, (1 - |z|^n)z)$ for all $z \in B^2$. You should prove in particular that f is well-defined, injective and surjective, and that both f and f^{-1} are continuous. You may assume that polynomials and the absolute value function are continuous, but beyond that you should not assume any properties of the given formula without proof. (13 marks)

- (b) For the boundary $S^1 \subset B^2$, explain briefly why S^1 / \sim is homeomorphic to S^1 again. (3 marks)
- (c) By adapting the method used for $\mathbb{R}P^2$, calculate $H_*(X)$. (14 marks)

(66) 2021-22 Mock Q5: Let X be a path connected space, and put

$$U = \{(t, x) \in S^1 \times X \mid t \neq (0, 1)\}$$
$$V = \{(t, x) \in S^1 \times X \mid t \neq (0, -1)\}$$

We use the usual notation for inclusion maps:

$$\begin{array}{ccc} U \cap V & & \stackrel{i}{\longrightarrow} & U \\ \downarrow & & & \downarrow k \\ V & & \stackrel{i}{\longrightarrow} & S^1 \times X \end{array}$$

- (a) Define maps $f, g: X \to U \cap V$ such that f gives a homotopy equivalence from X to one path component of $U \cap V$, and g gives a homotopy equivalence from X to the other path component of $U \cap V$. (4 marks)
- (b) Prove that the map $i' = i \circ f \colon X \to U \times X$ is homotopic to $i \circ g$, and also that i' is a homotopy equivalence. (You can then assume without further argument that the map $j' = j \circ f \colon X \to V \times X$ is homotopic to $j \circ g$, and that j' is a homotopy equivalence.) (6 marks)
- (c) Deduce descriptions of the homology groups $H_p(U \cap V)$, $H_p(U)$ and $H_p(V)$, and the homomorphism

$$\alpha = \begin{bmatrix} i_* \\ -j_* \end{bmatrix} \colon H_p(U \cap V) \to H_p(U) \oplus H_p(V).$$

Find the kernel and image of α . (8 marks)

- (d) Show that every element of $H_p(U) \oplus H_p(V)$ can be written as $(i'_*(a), 0) + \alpha(b)$ for a unique pair $(a, b) \in H_p(X)^2$. (3 marks)
- (e) Deduce that there is a short exact sequence $H_p(X) \to H_p(S^1 \times X) \to H_{p-1}(X)$. (4 marks)

(67) 2021-22 Q5: Put $X = \{(x, y) \in \mathbb{C}^2 \mid |x|^2 + |y|^2 = 1\}$, so X is homeomorphic to S^3 . Put $\omega = e^{2\pi i/3} \in \mathbb{C}$, so $\omega^3 = 1$. Define an equivalence relation on X by $(x, y) \sim (x', y')$ iff $(x', y') = \omega^k(x, y)$ for some k. Put

$$\begin{split} Y &= X / \sim \\ U &= \{ [x, y] \in Y \mid x \neq 0 \} \\ V &= \{ [x, y] \in Y \mid y \neq 0 \}. \end{split}$$

You may assume that U and V are open in Y and that $Y = U \cup V$.

- (a) Show that the formula $f([x, y]) = (x^3/|x|^3, y/x)$ gives a well-defined and continuous map $f: U \to S^1 \times \mathbb{C}$. Do not assume any properties of the given formula without checking them. (6 marks)
- (b) Show that f is actually a bijection and that the inverse satisfies

$$f^{-1}(u,z) = \left[(v,zv)/\sqrt{1+|z|^2} \right]$$

where v is any one of the three cube roots of u. Do not assume any properties of the given formula without checking them. (6 marks)

- (c) You may assume without proof that the map $f^{-1}: S^1 \times \mathbb{C} \to U$ is also continuous, so f is a homeomorphism. What can you conclude about the homeomorphism type of $U \cap V$? (3 marks)
- (d) The facts proved for U have obvious counterparts for V; you can assume these without proof. Deduce descriptions of $H_*(U)$, $H_*(V)$ and $H_*(U \cap V)$. (5 marks)

(e) Use the Mayer-Vietoris sequence to compute $H_*(Y)$. You should be able to compute $H_k(Y)$ for k = 0 and $k \ge 3$. For k = 1, 2 you will need to determine a map in the Mayer-Vietoris sequence, which is possible but not so easy. If you cannot see how to do it then you should guess, and give an answer based on your guess. (5 marks)

(68) 2022-23 Q1

- (a) Explain the terms homeomorphism and homeomorphic. (3 marks)
- (b) Explain the terms *homotopy*, *homotopic* and *homotopy equivalent*, distinguishing carefully between them. (5 marks)
- (c) Consider the following spaces:

$$X_0 = \{ z \in \mathbb{C} \mid \operatorname{Re}(z) \notin \mathbb{Z} \}$$
$$X_1 = \{ z \in \mathbb{C} \mid \operatorname{Im}(z) \in \mathbb{Z} \}$$
$$X_2 = \{ z \in \mathbb{C} \mid z \notin \mathbb{Z} \}$$
$$X_3 = \{ z \in \mathbb{R} \mid z \notin \mathbb{Z} \}$$
$$X_4 = \{ z \in \mathbb{C} \mid |z| \in \mathbb{Z} \}.$$

- (i) Sketch all these spaces. (5 marks)
- (ii) For which pairs (i, j) is X_i homotopy equivalent to X_j ? Justify your answer briefly. In cases where X_i is homotopy equivalent to X_j you should explain why, and in cases where X_i is not homotopy equivalent to X_j , you should explain that as well. (6 marks)
- (iii) For which pairs (i, j) is X_i homeomorphic to X_j ? Justify your answer briefly. In cases where X_i is homeomorphic to X_j you should explain why, and in cases where X_i is not homeomorphic to X_j , you should explain that as well. (6 marks)
- (69) 2022-23 Q5: Consider S^1 as the unit circle in \mathbb{R}^2 as usual. Let X be a path connected space, and put

$$U = \{(t, x) \in S^1 \times X \mid t \neq (0, 1)\}$$
$$V = \{(t, x) \in S^1 \times X \mid t \neq (0, -1)\}$$

We use the usual notation for inclusion maps:

$$\begin{array}{ccc} U \cap V & & \stackrel{i}{\longrightarrow} & U \\ \downarrow & & & \downarrow k \\ V & & \stackrel{i}{\longrightarrow} & S^1 \times X. \end{array}$$

- (a) Define maps $f, g: X \to U \cap V$ such that f gives a homotopy equivalence from X to one path component of $U \cap V$, and g gives a homotopy equivalence from X to the other path component of $U \cap V$. (4 marks)
- (b) Prove that the map $i' = i \circ f : X \to U$ is homotopic to $i \circ g$, and also that i' is a homotopy equivalence. (You can then assume without further argument that the map $j' = j \circ f : X \to V$ is homotopic to $j \circ g$, and that j' is a homotopy equivalence.) (6 marks)
- (c) Deduce descriptions (in terms of $H_*(X)$) of the homology groups $H_p(U \cap V)$, $H_p(U)$ and $H_p(V)$, and the homomorphism

$$\alpha = \begin{bmatrix} i_* \\ -j_* \end{bmatrix} \colon H_p(U \cap V) \to H_p(U) \oplus H_p(V)$$

Find the kernel and image of α . (8 marks)

- (d) Show that every element of $H_p(U) \oplus H_p(V)$ can be written as $(i'_*(a), 0) + \alpha(b)$ for a unique pair $(a, b) \in H_p(X) \oplus H_p(X)$. (3 marks)
- (e) Deduce that there is a short exact sequence $H_p(X) \to H_p(S^1 \times X) \to H_{p-1}(X)$. (4 marks)