

Chromatic cohomology of finite groups 3

Neil Strickland

December 4, 2023

- ▶ Again let $V \rightarrow X$ be a complex vector bundle of dimension d .
- ▶ $\text{Flag}_k(V) = \{(a, W_0, \dots, W_k) \mid x \in X, W_i < W_{i+1} \leq V_a, \dim(W_i) = i\}$.
- ▶ For $0 \leq i < k$ we have a line bundle $(Q_i)_{(a, \underline{W})} = W_{i+1}/W_i$ and an Euler class $x_i = e(Q_i) \in E^0(\text{Flag}_k(V))$.
- ▶ We also have a bundle R_k over $\text{Flag}_k(V)$ with $(R_k)_{(a, \underline{W})} = V_a/W_k$ (so $\dim(R_k) = d - k$), and $\text{Flag}_{k+1}(V)$ is the projective bundle $P(R_k)$.
- ▶ By induction based on this: the set of monomials $x^\alpha = \prod_{i < k} x_i^{\alpha_i}$ with $0 \leq \alpha_i < d - i$ is a basis for $E^0(\text{Flag}_k(V))$ over $E^0(X)$.
- ▶ For the ring structure: put $g_k(t) = \prod_{i < k} (t - x_i) \in E^0(X)[x_0, \dots, x_{k-1}]$, then divide $f_V(t)$ by $g(t)$ with remainder to get $f_V(t) = g(t)q(t) + r(t)$ with $\deg(r(t)) < k$, then let I be the ideal generated by the coefficients of $r(t)$. We then have $E^0(\text{Flag}_k(V)) = E^0(X)[x_0, \dots, x_{k-1}]/I$ as rings.
- ▶ Let G be a group with $|G| = n$. The representation $\mathbb{C}[G]$ gives a bundle $V = EG \times_G \mathbb{C}[G]$ over BG and a space $\text{Flag}_n(V)$ with $E^0(\text{Flag}_n(V)) \simeq E^0(BG)^{n!}$.
- ▶ $\text{Flag}_n(V) = EG \times_G F$, where $F = \{\underline{W} \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$.
- ▶ **Key fact:** all stabiliser groups in F are abelian. Indeed, $\text{stab}_G(\underline{W})$ injects in the abelian group $\prod_{i=0}^{n-1} \text{Aut}(W_{i+1} \ominus W_i) = (\mathbb{C}^\times)^n$.

- ▶ Again let $V \rightarrow X$ be a complex vector bundle of dimension d .
- ▶ $\text{Flag}_k(V) = \{(a, W_0, \dots, W_k) \mid x \in X, W_i < W_{i+1} \leq V_a, \dim(W_i) = i\}$.
- ▶ For $0 \leq i < k$ we have a line bundle $(Q_i)_{(a, \underline{W})} = W_{i+1}/W_i$ and an Euler class $x_i = e(Q_i) \in E^0(\text{Flag}_k(V))$.
- ▶ We also have a bundle R_k over $\text{Flag}_k(V)$ with $(R_k)_{(a, \underline{W})} = V_a/W_k$ (so $\dim(R_k) = d - k$), and $\text{Flag}_{k+1}(V)$ is the projective bundle $P(R_k)$.
- ▶ By induction based on this: the set of monomials $x^\alpha = \prod_{i < k} x_i^{\alpha_i}$ with $0 \leq \alpha_i < d - i$ is a basis for $E^0(\text{Flag}_k(V))$ over $E^0(X)$.
- ▶ For the ring structure: put $g_k(t) = \prod_{i < k} (t - x_i) \in E^0(X)[x_0, \dots, x_{k-1}]$, then divide $f_V(t)$ by $g(t)$ with remainder to get $f_V(t) = g(t)q(t) + r(t)$ with $\deg(r(t)) < k$, then let I be the ideal generated by the coefficients of $r(t)$. We then have $E^0(\text{Flag}_k(V)) = E^0(X)[x_0, \dots, x_{k-1}]/I$ as rings.
- ▶ Let G be a group with $|G| = n$. The representation $\mathbb{C}[G]$ gives a bundle $V = EG \times_G \mathbb{C}[G]$ over BG and a space $\text{Flag}_n(V)$ with $E^0(\text{Flag}_n(V)) \simeq E^0(BG)^{n!}$.
- ▶ $\text{Flag}_n(V) = EG \times_G F$, where $F = \{\underline{W} \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$.
- ▶ **Key fact:** all stabiliser groups in F are abelian. Indeed, $\text{stab}_G(\underline{W})$ injects in the abelian group $\prod_{i=0}^{n-1} \text{Aut}(W_{i+1} \ominus W_i) = (\mathbb{C}^\times)^n$.

- ▶ Again let $V \rightarrow X$ be a complex vector bundle of dimension d .
- ▶ $\text{Flag}_k(V) = \{(a, W_0, \dots, W_k) \mid x \in X, W_i < W_{i+1} \leq V_a, \dim(W_i) = i\}$.
- ▶ For $0 \leq i < k$ we have a line bundle $(Q_i)_{(a, \underline{W})} = W_{i+1}/W_i$ and an Euler class $x_i = e(Q_i) \in E^0(\text{Flag}_k(V))$.
- ▶ We also have a bundle R_k over $\text{Flag}_k(V)$ with $(R_k)_{(a, \underline{W})} = V_a/W_k$ (so $\dim(R_k) = d - k$), and $\text{Flag}_{k+1}(V)$ is the projective bundle $P(R_k)$.
- ▶ By induction based on this: the set of monomials $x^\alpha = \prod_{i < k} x_i^{\alpha_i}$ with $0 \leq \alpha_i < d - i$ is a basis for $E^0(\text{Flag}_k(V))$ over $E^0(X)$.
- ▶ For the ring structure: put $g_k(t) = \prod_{i < k} (t - x_i) \in E^0(X)[x_0, \dots, x_{k-1}]$, then divide $f_V(t)$ by $g(t)$ with remainder to get $f_V(t) = g(t)q(t) + r(t)$ with $\deg(r(t)) < k$, then let I be the ideal generated by the coefficients of $r(t)$. We then have $E^0(\text{Flag}_k(V)) = E^0(X)[x_0, \dots, x_{k-1}]/I$ as rings.
- ▶ Let G be a group with $|G| = n$. The representation $\mathbb{C}[G]$ gives a bundle $V = EG \times_G \mathbb{C}[G]$ over BG and a space $\text{Flag}_n(V)$ with $E^0(\text{Flag}_n(V)) \simeq E^0(BG)^{n!}$.
- ▶ $\text{Flag}_n(V) = EG \times_G F$, where $F = \{\underline{W} \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$.
- ▶ **Key fact:** all stabiliser groups in F are abelian. Indeed, $\text{stab}_G(\underline{W})$ injects in the abelian group $\prod_{i=0}^{n-1} \text{Aut}(W_{i+1} \ominus W_i) = (\mathbb{C}^\times)^n$.

- ▶ Again let $V \rightarrow X$ be a complex vector bundle of dimension d .
- ▶ $\text{Flag}_k(V) = \{(a, W_0, \dots, W_k) \mid x \in X, W_i < W_{i+1} \leq V_a, \dim(W_i) = i\}$.
- ▶ For $0 \leq i < k$ we have a line bundle $(Q_i)_{(a, \underline{W})} = W_{i+1}/W_i$ and an Euler class $x_i = e(Q_i) \in E^0(\text{Flag}_k(V))$.
- ▶ We also have a bundle R_k over $\text{Flag}_k(V)$ with $(R_k)_{(a, \underline{W})} = V_a/W_k$ (so $\dim(R_k) = d - k$), and $\text{Flag}_{k+1}(V)$ is the projective bundle $P(R_k)$.
- ▶ By induction based on this: the set of monomials $x^\alpha = \prod_{i < k} x_i^{\alpha_i}$ with $0 \leq \alpha_i < d - i$ is a basis for $E^0(\text{Flag}_k(V))$ over $E^0(X)$.
- ▶ For the ring structure: put $g_k(t) = \prod_{i < k} (t - x_i) \in E^0(X)[x_0, \dots, x_{k-1}]$, then divide $f_V(t)$ by $g(t)$ with remainder to get $f_V(t) = g(t)q(t) + r(t)$ with $\deg(r(t)) < k$, then let I be the ideal generated by the coefficients of $r(t)$. We then have $E^0(\text{Flag}_k(V)) = E^0(X)[x_0, \dots, x_{k-1}]/I$ as rings.
- ▶ Let G be a group with $|G| = n$. The representation $\mathbb{C}[G]$ gives a bundle $V = EG \times_G \mathbb{C}[G]$ over BG and a space $\text{Flag}_n(V)$ with $E^0(\text{Flag}_n(V)) \simeq E^0(BG)^{n!}$.
- ▶ $\text{Flag}_n(V) = EG \times_G F$, where $F = \{\underline{W} \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$.
- ▶ **Key fact:** all stabiliser groups in F are abelian. Indeed, $\text{stab}_G(\underline{W})$ injects in the abelian group $\prod_{i=0}^{n-1} \text{Aut}(W_{i+1} \ominus W_i) = (\mathbb{C}^\times)^n$.

- ▶ Again let $V \rightarrow X$ be a complex vector bundle of dimension d .
- ▶ $\text{Flag}_k(V) = \{(a, W_0, \dots, W_k) \mid x \in X, W_i < W_{i+1} \leq V_a, \dim(W_i) = i\}$.
- ▶ For $0 \leq i < k$ we have a line bundle $(Q_i)_{(a, \underline{W})} = W_{i+1}/W_i$ and an Euler class $x_i = e(Q_i) \in E^0(\text{Flag}_k(V))$.
- ▶ We also have a bundle R_k over $\text{Flag}_k(V)$ with $(R_k)_{(a, \underline{W})} = V_a/W_k$ (so $\dim(R_k) = d - k$), and $\text{Flag}_{k+1}(V)$ is the projective bundle $P(R_k)$.
- ▶ By induction based on this: the set of monomials $x^\alpha = \prod_{i < k} x_i^{\alpha_i}$ with $0 \leq \alpha_i < d - i$ is a basis for $E^0(\text{Flag}_k(V))$ over $E^0(X)$.
- ▶ For the ring structure: put $g_k(t) = \prod_{i < k} (t - x_i) \in E^0(X)[x_0, \dots, x_{k-1}]$, then divide $f_V(t)$ by $g(t)$ with remainder to get $f_V(t) = g(t)q(t) + r(t)$ with $\deg(r(t)) < k$, then let I be the ideal generated by the coefficients of $r(t)$. We then have $E^0(\text{Flag}_k(V)) = E^0(X)[x_0, \dots, x_{k-1}]/I$ as rings.
- ▶ Let G be a group with $|G| = n$. The representation $\mathbb{C}[G]$ gives a bundle $V = EG \times_G \mathbb{C}[G]$ over BG and a space $\text{Flag}_n(V)$ with $E^0(\text{Flag}_n(V)) \simeq E^0(BG)^{n!}$.
- ▶ $\text{Flag}_n(V) = EG \times_G F$, where $F = \{\underline{W} \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$.
- ▶ **Key fact:** all stabiliser groups in F are abelian. Indeed, $\text{stab}_G(\underline{W})$ injects in the abelian group $\prod_{i=0}^{n-1} \text{Aut}(W_{i+1} \ominus W_i) = (\mathbb{C}^\times)^n$.

- ▶ Again let $V \rightarrow X$ be a complex vector bundle of dimension d .
- ▶ $\text{Flag}_k(V) = \{(a, W_0, \dots, W_k) \mid x \in X, W_i < W_{i+1} \leq V_a, \dim(W_i) = i\}$.
- ▶ For $0 \leq i < k$ we have a line bundle $(Q_i)_{(a, \underline{W})} = W_{i+1}/W_i$ and an Euler class $x_i = e(Q_i) \in E^0(\text{Flag}_k(V))$.
- ▶ We also have a bundle R_k over $\text{Flag}_k(V)$ with $(R_k)_{(a, \underline{W})} = V_a/W_k$ (so $\dim(R_k) = d - k$), and $\text{Flag}_{k+1}(V)$ is the projective bundle $P(R_k)$.
- ▶ By induction based on this: the set of monomials $x^\alpha = \prod_{i < k} x_i^{\alpha_i}$ with $0 \leq \alpha_i < d - i$ is a basis for $E^0(\text{Flag}_k(V))$ over $E^0(X)$.
- ▶ For the ring structure: put $g_k(t) = \prod_{i < k} (t - x_i) \in E^0(X)[x_0, \dots, x_{k-1}]$, then divide $f_V(t)$ by $g(t)$ with remainder to get $f_V(t) = g(t)q(t) + r(t)$ with $\deg(r(t)) < k$, then let I be the ideal generated by the coefficients of $r(t)$. We then have $E^0(\text{Flag}_k(V)) = E^0(X)[x_0, \dots, x_{k-1}]/I$ as rings.
- ▶ Let G be a group with $|G| = n$. The representation $\mathbb{C}[G]$ gives a bundle $V = EG \times_G \mathbb{C}[G]$ over BG and a space $\text{Flag}_n(V)$ with $E^0(\text{Flag}_n(V)) \simeq E^0(BG)^{n!}$.
- ▶ $\text{Flag}_n(V) = EG \times_G F$, where $F = \{\underline{W} \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$.
- ▶ **Key fact:** all stabiliser groups in F are abelian. Indeed, $\text{stab}_G(\underline{W})$ injects in the abelian group $\prod_{i=0}^{n-1} \text{Aut}(W_{i+1} \ominus W_i) = (\mathbb{C}^\times)^n$.

- ▶ Again let $V \rightarrow X$ be a complex vector bundle of dimension d .
- ▶ $\text{Flag}_k(V) = \{(a, W_0, \dots, W_k) \mid x \in X, W_i < W_{i+1} \leq V_a, \dim(W_i) = i\}$.
- ▶ For $0 \leq i < k$ we have a line bundle $(Q_i)_{(a, \underline{W})} = W_{i+1}/W_i$ and an Euler class $x_i = e(Q_i) \in E^0(\text{Flag}_k(V))$.
- ▶ We also have a bundle R_k over $\text{Flag}_k(V)$ with $(R_k)_{(a, \underline{W})} = V_a/W_k$ (so $\dim(R_k) = d - k$), and $\text{Flag}_{k+1}(V)$ is the projective bundle $P(R_k)$.
- ▶ By induction based on this: the set of monomials $x^\alpha = \prod_{i < k} x_i^{\alpha_i}$ with $0 \leq \alpha_i < d - i$ is a basis for $E^0(\text{Flag}_k(V))$ over $E^0(X)$.
- ▶ For the ring structure: put $g_k(t) = \prod_{i < k} (t - x_i) \in E^0(X)[x_0, \dots, x_{k-1}]$, then divide $f_V(t)$ by $g(t)$ with remainder to get $f_V(t) = g(t)q(t) + r(t)$ with $\deg(r(t)) < k$, then let I be the ideal generated by the coefficients of $r(t)$. We then have $E^0(\text{Flag}_k(V)) = E^0(X)[x_0, \dots, x_{k-1}]/I$ as rings.
- ▶ Let G be a group with $|G| = n$. The representation $\mathbb{C}[G]$ gives a bundle $V = EG \times_G \mathbb{C}[G]$ over BG and a space $\text{Flag}_n(V)$ with $E^0(\text{Flag}_n(V)) \simeq E^0(BG)^{n!}$.
- ▶ $\text{Flag}_n(V) = EG \times_G F$, where $F = \{\underline{W} \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$.
- ▶ **Key fact:** all stabiliser groups in F are abelian. Indeed, $\text{stab}_G(\underline{W})$ injects in the abelian group $\prod_{i=0}^{n-1} \text{Aut}(W_{i+1} \ominus W_i) = (\mathbb{C}^\times)^n$.

- ▶ Again let $V \rightarrow X$ be a complex vector bundle of dimension d .
- ▶ $\text{Flag}_k(V) = \{(a, W_0, \dots, W_k) \mid x \in X, W_i < W_{i+1} \leq V_a, \dim(W_i) = i\}$.
- ▶ For $0 \leq i < k$ we have a line bundle $(Q_i)_{(a, \underline{W})} = W_{i+1}/W_i$ and an Euler class $x_i = e(Q_i) \in E^0(\text{Flag}_k(V))$.
- ▶ We also have a bundle R_k over $\text{Flag}_k(V)$ with $(R_k)_{(a, \underline{W})} = V_a/W_k$ (so $\dim(R_k) = d - k$), and $\text{Flag}_{k+1}(V)$ is the projective bundle $P(R_k)$.
- ▶ By induction based on this: the set of monomials $x^\alpha = \prod_{i < k} x_i^{\alpha_i}$ with $0 \leq \alpha_i < d - i$ is a basis for $E^0(\text{Flag}_k(V))$ over $E^0(X)$.
- ▶ For the ring structure: put $g_k(t) = \prod_{i < k} (t - x_i) \in E^0(X)[x_0, \dots, x_{k-1}]$, then divide $f_V(t)$ by $g(t)$ with remainder to get $f_V(t) = g(t)q(t) + r(t)$ with $\deg(r(t)) < k$, then let I be the ideal generated by the coefficients of $r(t)$. We then have $E^0(\text{Flag}_k(V)) = E^0(X)[x_0, \dots, x_{k-1}]/I$ as rings.
- ▶ Let G be a group with $|G| = n$. The representation $\mathbb{C}[G]$ gives a bundle $V = EG \times_G \mathbb{C}[G]$ over BG and a space $\text{Flag}_n(V)$ with $E^0(\text{Flag}_n(V)) \simeq E^0(BG)^{n!}$.
- ▶ $\text{Flag}_n(V) = EG \times_G F$, where $F = \{\underline{W} \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$.
- ▶ **Key fact:** all stabiliser groups in F are abelian. Indeed, $\text{stab}_G(\underline{W})$ injects in the abelian group $\prod_{i=0}^{n-1} \text{Aut}(W_{i+1} \ominus W_i) = (\mathbb{C}^\times)^n$.

- ▶ Again let $V \rightarrow X$ be a complex vector bundle of dimension d .
- ▶ $\text{Flag}_k(V) = \{(a, W_0, \dots, W_k) \mid x \in X, W_i < W_{i+1} \leq V_a, \dim(W_i) = i\}$.
- ▶ For $0 \leq i < k$ we have a line bundle $(Q_i)_{(a, \underline{W})} = W_{i+1}/W_i$ and an Euler class $x_i = e(Q_i) \in E^0(\text{Flag}_k(V))$.
- ▶ We also have a bundle R_k over $\text{Flag}_k(V)$ with $(R_k)_{(a, \underline{W})} = V_a/W_k$ (so $\dim(R_k) = d - k$), and $\text{Flag}_{k+1}(V)$ is the projective bundle $P(R_k)$.
- ▶ By induction based on this: the set of monomials $x^\alpha = \prod_{i < k} x_i^{\alpha_i}$ with $0 \leq \alpha_i < d - i$ is a basis for $E^0(\text{Flag}_k(V))$ over $E^0(X)$.
- ▶ For the ring structure: put $g_k(t) = \prod_{i < k} (t - x_i) \in E^0(X)[x_0, \dots, x_{k-1}]$, then divide $f_V(t)$ by $g(t)$ with remainder to get $f_V(t) = g(t)q(t) + r(t)$ with $\deg(r(t)) < k$, then let I be the ideal generated by the coefficients of $r(t)$. We then have $E^0(\text{Flag}_k(V)) = E^0(X)[x_0, \dots, x_{k-1}]/I$ as rings.
- ▶ Let G be a group with $|G| = n$. The representation $\mathbb{C}[G]$ gives a bundle $V = EG \times_G \mathbb{C}[G]$ over BG and a space $\text{Flag}_n(V)$ with $E^0(\text{Flag}_n(V)) \simeq E^0(BG)^{n!}$.
- ▶ $\text{Flag}_n(V) = EG \times_G F$, where $F = \{\underline{W} \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$.
- ▶ **Key fact:** all stabiliser groups in F are abelian. Indeed, $\text{stab}_G(\underline{W})$ injects in the abelian group $\prod_{i=0}^{n-1} \text{Aut}(W_{i+1} \ominus W_i) = (\mathbb{C}^\times)^n$.

- ▶ Again let $V \rightarrow X$ be a complex vector bundle of dimension d .
- ▶ $\text{Flag}_k(V) = \{(a, W_0, \dots, W_k) \mid x \in X, W_i < W_{i+1} \leq V_a, \dim(W_i) = i\}$.
- ▶ For $0 \leq i < k$ we have a line bundle $(Q_i)_{(a, \underline{W})} = W_{i+1}/W_i$ and an Euler class $x_i = e(Q_i) \in E^0(\text{Flag}_k(V))$.
- ▶ We also have a bundle R_k over $\text{Flag}_k(V)$ with $(R_k)_{(a, \underline{W})} = V_a/W_k$ (so $\dim(R_k) = d - k$), and $\text{Flag}_{k+1}(V)$ is the projective bundle $P(R_k)$.
- ▶ By induction based on this: the set of monomials $x^\alpha = \prod_{i < k} x_i^{\alpha_i}$ with $0 \leq \alpha_i < d - i$ is a basis for $E^0(\text{Flag}_k(V))$ over $E^0(X)$.
- ▶ For the ring structure: put $g_k(t) = \prod_{i < k} (t - x_i) \in E^0(X)[x_0, \dots, x_{k-1}]$, then divide $f_V(t)$ by $g(t)$ with remainder to get $f_V(t) = g(t)q(t) + r(t)$ with $\deg(r(t)) < k$, then let I be the ideal generated by the coefficients of $r(t)$. We then have $E^0(\text{Flag}_k(V)) = E^0(X)[x_0, \dots, x_{k-1}]/I$ as rings.
- ▶ Let G be a group with $|G| = n$. The representation $\mathbb{C}[G]$ gives a bundle $V = EG \times_G \mathbb{C}[G]$ over BG and a space $\text{Flag}_n(V)$ with $E^0(\text{Flag}_n(V)) \simeq E^0(BG)^{n!}$.
- ▶ $\text{Flag}_n(V) = EG \times_G F$, where $F = \{\underline{W} \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$.
- ▶ **Key fact:** all stabiliser groups in F are abelian. Indeed, $\text{stab}_G(\underline{W})$ injects in the abelian group $\prod_{i=0}^{n-1} \text{Aut}(W_{i+1} \ominus W_i) = (\mathbb{C}^\times)^n$.

- ▶ Again let $V \rightarrow X$ be a complex vector bundle of dimension d .
- ▶ $\text{Flag}_k(V) = \{(a, W_0, \dots, W_k) \mid x \in X, W_i < W_{i+1} \leq V_a, \dim(W_i) = i\}$.
- ▶ For $0 \leq i < k$ we have a line bundle $(Q_i)_{(a, \underline{W})} = W_{i+1}/W_i$ and an Euler class $x_i = e(Q_i) \in E^0(\text{Flag}_k(V))$.
- ▶ We also have a bundle R_k over $\text{Flag}_k(V)$ with $(R_k)_{(a, \underline{W})} = V_a/W_k$ (so $\dim(R_k) = d - k$), and $\text{Flag}_{k+1}(V)$ is the projective bundle $P(R_k)$.
- ▶ By induction based on this: the set of monomials $x^\alpha = \prod_{i < k} x_i^{\alpha_i}$ with $0 \leq \alpha_i < d - i$ is a basis for $E^0(\text{Flag}_k(V))$ over $E^0(X)$.
- ▶ For the ring structure: put $g_k(t) = \prod_{i < k} (t - x_i) \in E^0(X)[x_0, \dots, x_{k-1}]$, then divide $f_V(t)$ by $g(t)$ with remainder to get $f_V(t) = g(t)q(t) + r(t)$ with $\deg(r(t)) < k$, then let I be the ideal generated by the coefficients of $r(t)$. We then have $E^0(\text{Flag}_k(V)) = E^0(X)[x_0, \dots, x_{k-1}]/I$ as rings.
- ▶ Let G be a group with $|G| = n$. The representation $\mathbb{C}[G]$ gives a bundle $V = EG \times_G \mathbb{C}[G]$ over BG and a space $\text{Flag}_n(V)$ with $E^0(\text{Flag}_n(V)) \simeq E^0(BG)^{n!}$.
- ▶ $\text{Flag}_n(V) = EG \times_G F$, where $F = \{\underline{W} \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$.
- ▶ Key fact: all stabiliser groups in F are abelian. Indeed, $\text{stab}_G(\underline{W})$ injects in the abelian group $\prod_{i=0}^{n-1} \text{Aut}(W_{i+1} \ominus W_i) = (\mathbb{C}^\times)^n$.

- ▶ Again let $V \rightarrow X$ be a complex vector bundle of dimension d .
- ▶ $\text{Flag}_k(V) = \{(a, W_0, \dots, W_k) \mid x \in X, W_i < W_{i+1} \leq V_a, \dim(W_i) = i\}$.
- ▶ For $0 \leq i < k$ we have a line bundle $(Q_i)_{(a, \underline{W})} = W_{i+1}/W_i$ and an Euler class $x_i = e(Q_i) \in E^0(\text{Flag}_k(V))$.
- ▶ We also have a bundle R_k over $\text{Flag}_k(V)$ with $(R_k)_{(a, \underline{W})} = V_a/W_k$ (so $\dim(R_k) = d - k$), and $\text{Flag}_{k+1}(V)$ is the projective bundle $P(R_k)$.
- ▶ By induction based on this: the set of monomials $x^\alpha = \prod_{i < k} x_i^{\alpha_i}$ with $0 \leq \alpha_i < d - i$ is a basis for $E^0(\text{Flag}_k(V))$ over $E^0(X)$.
- ▶ For the ring structure: put $g_k(t) = \prod_{i < k} (t - x_i) \in E^0(X)[x_0, \dots, x_{k-1}]$, then divide $f_V(t)$ by $g(t)$ with remainder to get $f_V(t) = g(t)q(t) + r(t)$ with $\deg(r(t)) < k$, then let I be the ideal generated by the coefficients of $r(t)$. We then have $E^0(\text{Flag}_k(V)) = E^0(X)[x_0, \dots, x_{k-1}]/I$ as rings.
- ▶ Let G be a group with $|G| = n$. The representation $\mathbb{C}[G]$ gives a bundle $V = EG \times_G \mathbb{C}[G]$ over BG and a space $\text{Flag}_n(V)$ with $E^0(\text{Flag}_n(V)) \simeq E^0(BG)^{n!}$.
- ▶ $\text{Flag}_n(V) = EG \times_G F$, where $F = \{\underline{W} \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$.
- ▶ Key fact: all stabiliser groups in F are abelian. Indeed, $\text{stab}_G(\underline{W})$ injects in the abelian group $\prod_{i=0}^{n-1} \text{Aut}(W_{i+1} \ominus W_i) = (\mathbb{C}^\times)^n$.

- ▶ Again let $V \rightarrow X$ be a complex vector bundle of dimension d .
- ▶ $\text{Flag}_k(V) = \{(a, W_0, \dots, W_k) \mid x \in X, W_i < W_{i+1} \leq V_a, \dim(W_i) = i\}$.
- ▶ For $0 \leq i < k$ we have a line bundle $(Q_i)_{(a, \underline{W})} = W_{i+1}/W_i$ and an Euler class $x_i = e(Q_i) \in E^0(\text{Flag}_k(V))$.
- ▶ We also have a bundle R_k over $\text{Flag}_k(V)$ with $(R_k)_{(a, \underline{W})} = V_a/W_k$ (so $\dim(R_k) = d - k$), and $\text{Flag}_{k+1}(V)$ is the projective bundle $P(R_k)$.
- ▶ By induction based on this: the set of monomials $x^\alpha = \prod_{i < k} x_i^{\alpha_i}$ with $0 \leq \alpha_i < d - i$ is a basis for $E^0(\text{Flag}_k(V))$ over $E^0(X)$.
- ▶ For the ring structure: put $g_k(t) = \prod_{i < k} (t - x_i) \in E^0(X)[x_0, \dots, x_{k-1}]$, then divide $f_V(t)$ by $g(t)$ with remainder to get $f_V(t) = g(t)q(t) + r(t)$ with $\deg(r(t)) < k$, then let I be the ideal generated by the coefficients of $r(t)$. We then have $E^0(\text{Flag}_k(V)) = E^0(X)[x_0, \dots, x_{k-1}]/I$ as rings.
- ▶ Let G be a group with $|G| = n$. The representation $\mathbb{C}[G]$ gives a bundle $V = EG \times_G \mathbb{C}[G]$ over BG and a space $\text{Flag}_n(V)$ with $E^0(\text{Flag}_n(V)) \simeq E^0(BG)^{n!}$.
- ▶ $\text{Flag}_n(V) = EG \times_G F$, where $F = \{\underline{W} \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$.
- ▶ **Key fact:** all stabiliser groups in F are abelian. Indeed, $\text{stab}_G(\underline{W})$ injects in the abelian group $\prod_{i=0}^{n-1} \text{Aut}(W_{i+1} \ominus W_i) = (\mathbb{C}^\times)^n$.

- ▶ Again let $V \rightarrow X$ be a complex vector bundle of dimension d .
- ▶ $\text{Flag}_k(V) = \{(a, W_0, \dots, W_k) \mid x \in X, W_i < W_{i+1} \leq V_a, \dim(W_i) = i\}$.
- ▶ For $0 \leq i < k$ we have a line bundle $(Q_i)_{(a, \underline{W})} = W_{i+1}/W_i$ and an Euler class $x_i = e(Q_i) \in E^0(\text{Flag}_k(V))$.
- ▶ We also have a bundle R_k over $\text{Flag}_k(V)$ with $(R_k)_{(a, \underline{W})} = V_a/W_k$ (so $\dim(R_k) = d - k$), and $\text{Flag}_{k+1}(V)$ is the projective bundle $P(R_k)$.
- ▶ By induction based on this: the set of monomials $x^\alpha = \prod_{i < k} x_i^{\alpha_i}$ with $0 \leq \alpha_i < d - i$ is a basis for $E^0(\text{Flag}_k(V))$ over $E^0(X)$.
- ▶ For the ring structure: put $g_k(t) = \prod_{i < k} (t - x_i) \in E^0(X)[x_0, \dots, x_{k-1}]$, then divide $f_V(t)$ by $g(t)$ with remainder to get $f_V(t) = g(t)q(t) + r(t)$ with $\deg(r(t)) < k$, then let I be the ideal generated by the coefficients of $r(t)$. We then have $E^0(\text{Flag}_k(V)) = E^0(X)[x_0, \dots, x_{k-1}]/I$ as rings.
- ▶ Let G be a group with $|G| = n$. The representation $\mathbb{C}[G]$ gives a bundle $V = EG \times_G \mathbb{C}[G]$ over BG and a space $\text{Flag}_n(V)$ with $E^0(\text{Flag}_n(V)) \simeq E^0(BG)^{n!}$.
- ▶ $\text{Flag}_n(V) = EG \times_G F$, where $F = \{\underline{W} \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$.
- ▶ **Key fact:** all stabiliser groups in F are abelian. Indeed, $\text{stab}_G(\underline{W})$ injects in the abelian group $\prod_{i=0}^{n-1} \text{Aut}(W_{i+1} \ominus W_i) = (\mathbb{C}^\times)^n$.

- ▶ **Theorem:** if X is a finite simplicial complex with simplicial G -action, then the ring $E^*(EG \times_G X) = E^*(X_{hG})$ is finitely generated as an E^* -module.
- ▶ **Proof:** First treat the case $\text{stab}_G(x)$ is abelian for all $x \in X$.
- ▶ If $X = G/H$ then H must be abelian and $X_{hG} = BH$ and $E^*(BH)$ is finitely generated by previous calculation.
- ▶ If X is just a finite discrete G -set then it is a disjoint union of G/H 's and the same applies.
- ▶ In general, if X^k is the k -skeleton of X then $X^k/X^{k-1} = \Sigma^k W_+$ for some finite G -set W , giving an exact sequence of $E^*(1)$ -modules

$$E^{*-k}(W_{hG}) \rightarrow E^*(X_{hG}^k) \rightarrow E^*(X_{hG}^{k-1}).$$

As $E^*(1)$ is noetherian, it follows inductively that $E^*(X_{hG}^k)$ is finitely generated for all k , so $E^*(X_{hG})$ is finitely generated.

- ▶ Now remove the abelian stabiliser condition.
- ▶ Put $n = |G|$ and $F = \{(W_0, \dots, W_n) \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$.
- ▶ Then $X \times F$ has abelian stabilisers, so $A^* = E^*((X \times F)_{hG})$ is finitely generated; enough to show that $B^* = E^*(X_{hG})$ is a retract of this.
- ▶ But $(X \times F)_{hG}$ is $\text{Flag}_n(V)$ for a bundle V over BG , so $A^* \simeq (B^*)^{n!}$, so B^* is a retract of A^* \square .

- ▶ **Theorem:** if X is a finite simplicial complex with simplicial G -action, then the ring $E^*(EG \times_G X) = E^*(X_{hG})$ is finitely generated as an E^* -module.
- ▶ **Proof:** First treat the case $\text{stab}_G(x)$ is abelian for all $x \in X$.
- ▶ If $X = G/H$ then H must be abelian and $X_{hG} = BH$ and $E^*(BH)$ is finitely generated by previous calculation.
- ▶ If X is just a finite discrete G -set then it is a disjoint union of G/H 's and the same applies.
- ▶ In general, if X^k is the k -skeleton of X then $X^k/X^{k-1} = \Sigma^k W_+$ for some finite G -set W , giving an exact sequence of $E^*(1)$ -modules

$$E^{*-k}(W_{hG}) \rightarrow E^*(X_{hG}^k) \rightarrow E^*(X_{hG}^{k-1}).$$

As $E^*(1)$ is noetherian, it follows inductively that $E^*(X_{hG}^k)$ is finitely generated for all k , so $E^*(X_{hG})$ is finitely generated.

- ▶ Now remove the abelian stabiliser condition.
- ▶ Put $n = |G|$ and $F = \{(W_0, \dots, W_n) \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$.
- ▶ Then $X \times F$ has abelian stabilisers, so $A^* = E^*((X \times F)_{hG})$ is finitely generated; enough to show that $B^* = E^*(X_{hG})$ is a retract of this.
- ▶ But $(X \times F)_{hG}$ is $\text{Flag}_n(V)$ for a bundle V over BG , so $A^* \simeq (B^*)^{n!}$, so B^* is a retract of A^* \square .

- ▶ **Theorem:** if X is a finite simplicial complex with simplicial G -action, then the ring $E^*(EG \times_G X) = E^*(X_{hG})$ is finitely generated as an E^* -module.
- ▶ **Proof:** First treat the case $\text{stab}_G(x)$ is abelian for all $x \in X$.
- ▶ If $X = G/H$ then H must be abelian and $X_{hG} = BH$ and $E^*(BH)$ is finitely generated by previous calculation.
- ▶ If X is just a finite discrete G -set then it is a disjoint union of G/H 's and the same applies.
- ▶ In general, if X^k is the k -skeleton of X then $X^k/X^{k-1} = \Sigma^k W_+$ for some finite G -set W , giving an exact sequence of $E^*(1)$ -modules

$$E^{*-k}(W_{hG}) \rightarrow E^*(X_{hG}^k) \rightarrow E^*(X_{hG}^{k-1}).$$

As $E^*(1)$ is noetherian, it follows inductively that $E^*(X_{hG}^k)$ is finitely generated for all k , so $E^*(X_{hG})$ is finitely generated.

- ▶ Now remove the abelian stabiliser condition.
- ▶ Put $n = |G|$ and $F = \{(W_0, \dots, W_n) \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$.
- ▶ Then $X \times F$ has abelian stabilisers, so $A^* = E^*((X \times F)_{hG})$ is finitely generated; enough to show that $B^* = E^*(X_{hG})$ is a retract of this.
- ▶ But $(X \times F)_{hG}$ is $\text{Flag}_n(V)$ for a bundle V over BG , so $A^* \simeq (B^*)^{n!}$, so B^* is a retract of A^* \square .

- ▶ **Theorem:** if X is a finite simplicial complex with simplicial G -action, then the ring $E^*(EG \times_G X) = E^*(X_{hG})$ is finitely generated as an E^* -module.
- ▶ **Proof:** First treat the case $\text{stab}_G(x)$ is abelian for all $x \in X$.
- ▶ If $X = G/H$ then H must be abelian and $X_{hG} = BH$ and $E^*(BH)$ is finitely generated by previous calculation.
- ▶ If X is just a finite discrete G -set then it is a disjoint union of G/H 's and the same applies.
- ▶ In general, if X^k is the k -skeleton of X then $X^k/X^{k-1} = \Sigma^k W_+$ for some finite G -set W , giving an exact sequence of $E^*(1)$ -modules

$$E^{*-k}(W_{hG}) \rightarrow E^*(X_{hG}^k) \rightarrow E^*(X_{hG}^{k-1}).$$

As $E^*(1)$ is noetherian, it follows inductively that $E^*(X_{hG}^k)$ is finitely generated for all k , so $E^*(X_{hG})$ is finitely generated.

- ▶ Now remove the abelian stabiliser condition.
- ▶ Put $n = |G|$ and $F = \{(W_0, \dots, W_n) \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$.
- ▶ Then $X \times F$ has abelian stabilisers, so $A^* = E^*((X \times F)_{hG})$ is finitely generated; enough to show that $B^* = E^*(X_{hG})$ is a retract of this.
- ▶ But $(X \times F)_{hG}$ is $\text{Flag}_n(V)$ for a bundle V over BG , so $A^* \simeq (B^*)^{n!}$, so B^* is a retract of A^* \square .

- ▶ **Theorem:** if X is a finite simplicial complex with simplicial G -action, then the ring $E^*(EG \times_G X) = E^*(X_{hG})$ is finitely generated as an E^* -module.
- ▶ **Proof:** First treat the case $\text{stab}_G(x)$ is abelian for all $x \in X$.
- ▶ If $X = G/H$ then H must be abelian and $X_{hG} = BH$ and $E^*(BH)$ is finitely generated by previous calculation.
- ▶ If X is just a finite discrete G -set then it is a disjoint union of G/H 's and the same applies.
- ▶ In general, if X^k is the k -skeleton of X then $X^k/X^{k-1} = \Sigma^k W_+$ for some finite G -set W , giving an exact sequence of $E^*(1)$ -modules

$$E^{*-k}(W_{hG}) \rightarrow E^*(X_{hG}^k) \rightarrow E^*(X_{hG}^{k-1}).$$

As $E^*(1)$ is noetherian, it follows inductively that $E^*(X_{hG}^k)$ is finitely generated for all k , so $E^*(X_{hG})$ is finitely generated.

- ▶ Now remove the abelian stabiliser condition.
- ▶ Put $n = |G|$ and $F = \{(W_0, \dots, W_n) \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$.
- ▶ Then $X \times F$ has abelian stabilisers, so $A^* = E^*((X \times F)_{hG})$ is finitely generated; enough to show that $B^* = E^*(X_{hG})$ is a retract of this.
- ▶ But $(X \times F)_{hG}$ is $\text{Flag}_n(V)$ for a bundle V over BG , so $A^* \simeq (B^*)^{n!}$, so B^* is a retract of A^* \square .

- ▶ **Theorem:** if X is a finite simplicial complex with simplicial G -action, then the ring $E^*(EG \times_G X) = E^*(X_{hG})$ is finitely generated as an E^* -module.
- ▶ **Proof:** First treat the case $\text{stab}_G(x)$ is abelian for all $x \in X$.
- ▶ If $X = G/H$ then H must be abelian and $X_{hG} = BH$ and $E^*(BH)$ is finitely generated by previous calculation.
- ▶ If X is just a finite discrete G -set then it is a disjoint union of G/H 's and the same applies.
- ▶ In general, if X^k is the k -skeleton of X then $X^k/X^{k-1} = \Sigma^k W_+$ for some finite G -set W , giving an exact sequence of $E^*(1)$ -modules

$$E^{*-k}(W_{hG}) \rightarrow E^*(X_{hG}^k) \rightarrow E^*(X_{hG}^{k-1}).$$

As $E^*(1)$ is noetherian, it follows inductively that $E^*(X_{hG}^k)$ is finitely generated for all k , so $E^*(X_{hG})$ is finitely generated.

- ▶ Now remove the abelian stabiliser condition.
- ▶ Put $n = |G|$ and $F = \{(W_0, \dots, W_n) \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$.
- ▶ Then $X \times F$ has abelian stabilisers, so $A^* = E^*((X \times F)_{hG})$ is finitely generated; enough to show that $B^* = E^*(X_{hG})$ is a retract of this.
- ▶ But $(X \times F)_{hG}$ is $\text{Flag}_n(V)$ for a bundle V over BG , so $A^* \simeq (B^*)^{n!}$, so B^* is a retract of A^* \square .

- ▶ **Theorem:** if X is a finite simplicial complex with simplicial G -action, then the ring $E^*(EG \times_G X) = E^*(X_{hG})$ is finitely generated as an E^* -module.
- ▶ **Proof:** First treat the case $\text{stab}_G(x)$ is abelian for all $x \in X$.
- ▶ If $X = G/H$ then H must be abelian and $X_{hG} = BH$ and $E^*(BH)$ is finitely generated by previous calculation.
- ▶ If X is just a finite discrete G -set then it is a disjoint union of G/H 's and the same applies.
- ▶ In general, if X^k is the k -skeleton of X then $X^k/X^{k-1} = \Sigma^k W_+$ for some finite G -set W , giving an exact sequence of $E^*(1)$ -modules

$$E^{*-k}(W_{hG}) \rightarrow E^*(X_{hG}^k) \rightarrow E^*(X_{hG}^{k-1}).$$

As $E^*(1)$ is noetherian, it follows inductively that $E^*(X_{hG}^k)$ is finitely generated for all k , so $E^*(X_{hG})$ is finitely generated.

- ▶ Now remove the abelian stabiliser condition.
- ▶ Put $n = |G|$ and $F = \{(W_0, \dots, W_n) \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$.
- ▶ Then $X \times F$ has abelian stabilisers, so $A^* = E^*((X \times F)_{hG})$ is finitely generated; enough to show that $B^* = E^*(X_{hG})$ is a retract of this.
- ▶ But $(X \times F)_{hG}$ is $\text{Flag}_n(V)$ for a bundle V over BG , so $A^* \simeq (B^*)^{n!}$, so B^* is a retract of A^* \square .

- ▶ **Theorem:** if X is a finite simplicial complex with simplicial G -action, then the ring $E^*(EG \times_G X) = E^*(X_{hG})$ is finitely generated as an E^* -module.
- ▶ **Proof:** First treat the case $\text{stab}_G(x)$ is abelian for all $x \in X$.
- ▶ If $X = G/H$ then H must be abelian and $X_{hG} = BH$ and $E^*(BH)$ is finitely generated by previous calculation.
- ▶ If X is just a finite discrete G -set then it is a disjoint union of G/H 's and the same applies.
- ▶ In general, if X^k is the k -skeleton of X then $X^k/X^{k-1} = \Sigma^k W_+$ for some finite G -set W , giving an exact sequence of $E^*(1)$ -modules

$$E^{*-k}(W_{hG}) \rightarrow E^*(X_{hG}^k) \rightarrow E^*(X_{hG}^{k-1}).$$

As $E^*(1)$ is noetherian, it follows inductively that $E^*(X_{hG}^k)$ is finitely generated for all k , so $E^*(X_{hG})$ is finitely generated.

- ▶ Now remove the abelian stabiliser condition.
- ▶ Put $n = |G|$ and $F = \{(W_0, \dots, W_n) \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$.
- ▶ Then $X \times F$ has abelian stabilisers, so $A^* = E^*((X \times F)_{hG})$ is finitely generated; enough to show that $B^* = E^*(X_{hG})$ is a retract of this.
- ▶ But $(X \times F)_{hG}$ is $\text{Flag}_n(V)$ for a bundle V over BG , so $A^* \simeq (B^*)^{n!}$, so B^* is a retract of A^* \square .

- ▶ **Theorem:** if X is a finite simplicial complex with simplicial G -action, then the ring $E^*(EG \times_G X) = E^*(X_{hG})$ is finitely generated as an E^* -module.
- ▶ **Proof:** First treat the case $\text{stab}_G(x)$ is abelian for all $x \in X$.
- ▶ If $X = G/H$ then H must be abelian and $X_{hG} = BH$ and $E^*(BH)$ is finitely generated by previous calculation.
- ▶ If X is just a finite discrete G -set then it is a disjoint union of G/H 's and the same applies.
- ▶ In general, if X^k is the k -skeleton of X then $X^k/X^{k-1} = \Sigma^k W_+$ for some finite G -set W , giving an exact sequence of $E^*(1)$ -modules

$$E^{*-k}(W_{hG}) \rightarrow E^*(X_{hG}^k) \rightarrow E^*(X_{hG}^{k-1}).$$

As $E^*(1)$ is noetherian, it follows inductively that $E^*(X_{hG}^k)$ is finitely generated for all k , so $E^*(X_{hG})$ is finitely generated.

- ▶ Now remove the abelian stabiliser condition.
- ▶ Put $n = |G|$ and $F = \{(W_0, \dots, W_n) \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$.
- ▶ Then $X \times F$ has abelian stabilisers, so $A^* = E^*((X \times F)_{hG})$ is finitely generated; enough to show that $B^* = E^*(X_{hG})$ is a retract of this.
- ▶ But $(X \times F)_{hG}$ is $\text{Flag}_n(V)$ for a bundle V over BG , so $A^* \simeq (B^*)^{n!}$, so B^* is a retract of A^* \square .

- ▶ **Theorem:** if X is a finite simplicial complex with simplicial G -action, then the ring $E^*(EG \times_G X) = E^*(X_{hG})$ is finitely generated as an E^* -module.
- ▶ **Proof:** First treat the case $\text{stab}_G(x)$ is abelian for all $x \in X$.
- ▶ If $X = G/H$ then H must be abelian and $X_{hG} = BH$ and $E^*(BH)$ is finitely generated by previous calculation.
- ▶ If X is just a finite discrete G -set then it is a disjoint union of G/H 's and the same applies.
- ▶ In general, if X^k is the k -skeleton of X then $X^k/X^{k-1} = \Sigma^k W_+$ for some finite G -set W , giving an exact sequence of $E^*(1)$ -modules

$$E^{*-k}(W_{hG}) \rightarrow E^*(X_{hG}^k) \rightarrow E^*(X_{hG}^{k-1}).$$

As $E^*(1)$ is noetherian, it follows inductively that $E^*(X_{hG}^k)$ is finitely generated for all k , so $E^*(X_{hG})$ is finitely generated.

- ▶ Now remove the abelian stabiliser condition.
- ▶ Put $n = |G|$ and $F = \{(W_0, \dots, W_n) \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$.
- ▶ Then $X \times F$ has abelian stabilisers, so $A^* = E^*((X \times F)_{hG})$ is finitely generated; enough to show that $B^* = E^*(X_{hG})$ is a retract of this.
- ▶ But $(X \times F)_{hG}$ is $\text{Flag}_n(V)$ for a bundle V over BG , so $A^* \simeq (B^*)^{n!}$, so B^* is a retract of A^* \square .

- ▶ **Theorem:** if X is a finite simplicial complex with simplicial G -action, then the ring $E^*(EG \times_G X) = E^*(X_{hG})$ is finitely generated as an E^* -module.
- ▶ **Proof:** First treat the case $\text{stab}_G(x)$ is abelian for all $x \in X$.
- ▶ If $X = G/H$ then H must be abelian and $X_{hG} = BH$ and $E^*(BH)$ is finitely generated by previous calculation.
- ▶ If X is just a finite discrete G -set then it is a disjoint union of G/H 's and the same applies.
- ▶ In general, if X^k is the k -skeleton of X then $X^k/X^{k-1} = \Sigma^k W_+$ for some finite G -set W , giving an exact sequence of $E^*(1)$ -modules

$$E^{*-k}(W_{hG}) \rightarrow E^*(X_{hG}^k) \rightarrow E^*(X_{hG}^{k-1}).$$

As $E^*(1)$ is noetherian, it follows inductively that $E^*(X_{hG}^k)$ is finitely generated for all k , so $E^*(X_{hG})$ is finitely generated.

- ▶ Now remove the abelian stabiliser condition.
- ▶ Put $n = |G|$ and $F = \{(W_0, \dots, W_n) \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$.
- ▶ Then $X \times F$ has abelian stabilisers, so $A^* = E^*((X \times F)_{hG})$ is finitely generated; enough to show that $B^* = E^*(X_{hG})$ is a retract of this.
- ▶ But $(X \times F)_{hG}$ is $\text{Flag}_n(V)$ for a bundle V over BG , so $A^* \simeq (B^*)^{n!}$, so B^* is a retract of A^* \square .

Finite groupoids

- ▶ Many things about Morava E -theory are more convenient using groupoids.
- ▶ A *groupoid* is a category G in which all morphisms are invertible.
- ▶ Say G is *finite* if all Hom sets $G(a, b)$ are finite, and the set $\pi_0(G)$ of isomorphism classes is finite.
- ▶ If so, we can choose a_1, \dots, a_m containing one element of each isomorphism class, and put $G_i = G(a_i, a_i)$, and we get $BG \simeq \coprod_i BG_i$.
- ▶ Thus $E^*(BG) = \prod_i E^*(BG_i)$, which is a finitely generated E^* -module.
- ▶ Any group can be regarded as a groupoid with one object.
- ▶ A *representation* of G is a functor V from G to the category \mathcal{V} of finite-dimensional complex vector spaces.
- ▶ This again gives spaces $\text{Flag}_k(V)$ and $P(V) = \text{Flag}_1(V)$ over BG .
- ▶ Given groupoids G and H , the functor category $[G, H]$ is also a groupoid.
- ▶ If G, H are groups then $\text{obj}([G, H]) = \text{Hom}(G, H)$ and morphisms $\alpha \rightarrow \beta$ in $[G, H]$ are elements $h \in H$ with $\beta(g) = h\alpha(g)h^{-1}$ for all $g \in G$.
- ▶ So $\alpha \simeq \beta$ iff α and β are conjugate, and $\pi_0([G, H])$ is the set of conjugacy classes of homomorphisms.

- ▶ Many things about Morava E -theory are more convenient using groupoids.
- ▶ A *groupoid* is a category G in which all morphisms are invertible.
- ▶ Say G is *finite* if all Hom sets $G(a, b)$ are finite, and the set $\pi_0(G)$ of isomorphism classes is finite.
- ▶ If so, we can choose a_1, \dots, a_m containing one element of each isomorphism class, and put $G_i = G(a_i, a_i)$, and we get $BG \simeq \coprod_i BG_i$.
- ▶ Thus $E^*(BG) = \prod_i E^*(BG_i)$, which is a finitely generated E^* -module.
- ▶ Any group can be regarded as a groupoid with one object.
- ▶ A *representation* of G is a functor V from G to the category \mathcal{V} of finite-dimensional complex vector spaces.
- ▶ This again gives spaces $\text{Flag}_k(V)$ and $P(V) = \text{Flag}_1(V)$ over BG .
- ▶ Given groupoids G and H , the functor category $[G, H]$ is also a groupoid.
- ▶ If G, H are groups then $\text{obj}([G, H]) = \text{Hom}(G, H)$ and morphisms $\alpha \rightarrow \beta$ in $[G, H]$ are elements $h \in H$ with $\beta(g) = h\alpha(g)h^{-1}$ for all $g \in G$.
- ▶ So $\alpha \simeq \beta$ iff α and β are conjugate, and $\pi_0([G, H])$ is the set of conjugacy classes of homomorphisms.

Finite groupoids

- ▶ Many things about Morava E -theory are more convenient using groupoids.
- ▶ A *groupoid* is a category G in which all morphisms are invertible.
- ▶ Say G is *finite* if all Hom sets $G(a, b)$ are finite, and the set $\pi_0(G)$ of isomorphism classes is finite.
- ▶ If so, we can choose a_1, \dots, a_m containing one element of each isomorphism class, and put $G_i = G(a_i, a_i)$, and we get $BG \simeq \coprod_i BG_i$.
- ▶ Thus $E^*(BG) = \prod_i E^*(BG_i)$, which is a finitely generated E^* -module.
- ▶ Any group can be regarded as a groupoid with one object.
- ▶ A *representation* of G is a functor V from G to the category \mathcal{V} of finite-dimensional complex vector spaces.
- ▶ This again gives spaces $\text{Flag}_k(V)$ and $P(V) = \text{Flag}_1(V)$ over BG .
- ▶ Given groupoids G and H , the functor category $[G, H]$ is also a groupoid.
- ▶ If G, H are groups then $\text{obj}([G, H]) = \text{Hom}(G, H)$ and morphisms $\alpha \rightarrow \beta$ in $[G, H]$ are elements $h \in H$ with $\beta(g) = h\alpha(g)h^{-1}$ for all $g \in G$.
- ▶ So $\alpha \simeq \beta$ iff α and β are conjugate, and $\pi_0([G, H])$ is the set of conjugacy classes of homomorphisms.

- ▶ Many things about Morava E -theory are more convenient using groupoids.
- ▶ A *groupoid* is a category G in which all morphisms are invertible.
- ▶ Say G is *finite* if all Hom sets $G(a, b)$ are finite, and the set $\pi_0(G)$ of isomorphism classes is finite.
- ▶ If so, we can choose a_1, \dots, a_m containing one element of each isomorphism class, and put $G_i = G(a_i, a_i)$, and we get $BG \simeq \coprod_i BG_i$.
- ▶ Thus $E^*(BG) = \coprod_i E^*(BG_i)$, which is a finitely generated E^* -module.
- ▶ Any group can be regarded as a groupoid with one object.
- ▶ A *representation* of G is a functor V from G to the category \mathcal{V} of finite-dimensional complex vector spaces.
- ▶ This again gives spaces $\text{Flag}_k(V)$ and $P(V) = \text{Flag}_1(V)$ over BG .
- ▶ Given groupoids G and H , the functor category $[G, H]$ is also a groupoid.
- ▶ If G, H are groups then $\text{obj}([G, H]) = \text{Hom}(G, H)$ and morphisms $\alpha \rightarrow \beta$ in $[G, H]$ are elements $h \in H$ with $\beta(g) = h\alpha(g)h^{-1}$ for all $g \in G$.
- ▶ So $\alpha \simeq \beta$ iff α and β are conjugate, and $\pi_0([G, H])$ is the set of conjugacy classes of homomorphisms.

- ▶ Many things about Morava E -theory are more convenient using groupoids.
- ▶ A *groupoid* is a category G in which all morphisms are invertible.
- ▶ Say G is *finite* if all Hom sets $G(a, b)$ are finite, and the set $\pi_0(G)$ of isomorphism classes is finite.
- ▶ If so, we can choose a_1, \dots, a_m containing one element of each isomorphism class, and put $G_i = G(a_i, a_i)$, and we get $BG \simeq \coprod_i BG_i$.
- ▶ Thus $E^*(BG) = \coprod_i E^*(BG_i)$, which is a finitely generated E^* -module.
- ▶ Any group can be regarded as a groupoid with one object.
- ▶ A *representation* of G is a functor V from G to the category \mathcal{V} of finite-dimensional complex vector spaces.
- ▶ This again gives spaces $\text{Flag}_k(V)$ and $P(V) = \text{Flag}_1(V)$ over BG .
- ▶ Given groupoids G and H , the functor category $[G, H]$ is also a groupoid.
- ▶ If G, H are groups then $\text{obj}([G, H]) = \text{Hom}(G, H)$ and morphisms $\alpha \rightarrow \beta$ in $[G, H]$ are elements $h \in H$ with $\beta(g) = h\alpha(g)h^{-1}$ for all $g \in G$.
- ▶ So $\alpha \simeq \beta$ iff α and β are conjugate, and $\pi_0([G, H])$ is the set of conjugacy classes of homomorphisms.

- ▶ Many things about Morava E -theory are more convenient using groupoids.
- ▶ A *groupoid* is a category G in which all morphisms are invertible.
- ▶ Say G is *finite* if all Hom sets $G(a, b)$ are finite, and the set $\pi_0(G)$ of isomorphism classes is finite.
- ▶ If so, we can choose a_1, \dots, a_m containing one element of each isomorphism class, and put $G_i = G(a_i, a_i)$, and we get $BG \simeq \coprod_i BG_i$.
- ▶ Thus $E^*(BG) = \prod_i E^*(BG_i)$, which is a finitely generated E^* -module.
- ▶ Any group can be regarded as a groupoid with one object.
- ▶ A *representation* of G is a functor V from G to the category \mathcal{V} of finite-dimensional complex vector spaces.
- ▶ This again gives spaces $\text{Flag}_k(V)$ and $P(V) = \text{Flag}_1(V)$ over BG .
- ▶ Given groupoids G and H , the functor category $[G, H]$ is also a groupoid.
- ▶ If G, H are groups then $\text{obj}([G, H]) = \text{Hom}(G, H)$ and morphisms $\alpha \rightarrow \beta$ in $[G, H]$ are elements $h \in H$ with $\beta(g) = h\alpha(g)h^{-1}$ for all $g \in G$.
- ▶ So $\alpha \simeq \beta$ iff α and β are conjugate, and $\pi_0([G, H])$ is the set of conjugacy classes of homomorphisms.

- ▶ Many things about Morava E -theory are more convenient using groupoids.
- ▶ A *groupoid* is a category G in which all morphisms are invertible.
- ▶ Say G is *finite* if all Hom sets $G(a, b)$ are finite, and the set $\pi_0(G)$ of isomorphism classes is finite.
- ▶ If so, we can choose a_1, \dots, a_m containing one element of each isomorphism class, and put $G_i = G(a_i, a_i)$, and we get $BG \simeq \coprod_i BG_i$.
- ▶ Thus $E^*(BG) = \prod_i E^*(BG_i)$, which is a finitely generated E^* -module.
- ▶ Any group can be regarded as a groupoid with one object.
- ▶ A *representation* of G is a functor V from G to the category \mathcal{V} of finite-dimensional complex vector spaces.
- ▶ This again gives spaces $\text{Flag}_k(V)$ and $P(V) = \text{Flag}_1(V)$ over BG .
- ▶ Given groupoids G and H , the functor category $[G, H]$ is also a groupoid.
- ▶ If G, H are groups then $\text{obj}([G, H]) = \text{Hom}(G, H)$ and morphisms $\alpha \rightarrow \beta$ in $[G, H]$ are elements $h \in H$ with $\beta(g) = h\alpha(g)h^{-1}$ for all $g \in G$.
- ▶ So $\alpha \simeq \beta$ iff α and β are conjugate, and $\pi_0([G, H])$ is the set of conjugacy classes of homomorphisms.

- ▶ Many things about Morava E -theory are more convenient using groupoids.
- ▶ A *groupoid* is a category G in which all morphisms are invertible.
- ▶ Say G is *finite* if all Hom sets $G(a, b)$ are finite, and the set $\pi_0(G)$ of isomorphism classes is finite.
- ▶ If so, we can choose a_1, \dots, a_m containing one element of each isomorphism class, and put $G_i = G(a_i, a_i)$, and we get $BG \simeq \coprod_i BG_i$.
- ▶ Thus $E^*(BG) = \prod_i E^*(BG_i)$, which is a finitely generated E^* -module.
- ▶ Any group can be regarded as a groupoid with one object.
- ▶ A *representation* of G is a functor V from G to the category \mathcal{V} of finite-dimensional complex vector spaces.
 - ▶ This again gives spaces $\text{Flag}_k(V)$ and $P(V) = \text{Flag}_1(V)$ over BG .
 - ▶ Given groupoids G and H , the functor category $[G, H]$ is also a groupoid.
 - ▶ If G, H are groups then $\text{obj}([G, H]) = \text{Hom}(G, H)$ and morphisms $\alpha \rightarrow \beta$ in $[G, H]$ are elements $h \in H$ with $\beta(g) = h\alpha(g)h^{-1}$ for all $g \in G$.
 - ▶ So $\alpha \simeq \beta$ iff α and β are conjugate, and $\pi_0([G, H])$ is the set of conjugacy classes of homomorphisms.

- ▶ Many things about Morava E -theory are more convenient using groupoids.
- ▶ A *groupoid* is a category G in which all morphisms are invertible.
- ▶ Say G is *finite* if all Hom sets $G(a, b)$ are finite, and the set $\pi_0(G)$ of isomorphism classes is finite.
- ▶ If so, we can choose a_1, \dots, a_m containing one element of each isomorphism class, and put $G_i = G(a_i, a_i)$, and we get $BG \simeq \coprod_i BG_i$.
- ▶ Thus $E^*(BG) = \prod_i E^*(BG_i)$, which is a finitely generated E^* -module.
- ▶ Any group can be regarded as a groupoid with one object.
- ▶ A *representation* of G is a functor V from G to the category \mathcal{V} of finite-dimensional complex vector spaces.
- ▶ This again gives spaces $\text{Flag}_k(V)$ and $P(V) = \text{Flag}_1(V)$ over BG .
- ▶ Given groupoids G and H , the functor category $[G, H]$ is also a groupoid.
- ▶ If G, H are groups then $\text{obj}([G, H]) = \text{Hom}(G, H)$ and morphisms $\alpha \rightarrow \beta$ in $[G, H]$ are elements $h \in H$ with $\beta(g) = h\alpha(g)h^{-1}$ for all $g \in G$.
- ▶ So $\alpha \simeq \beta$ iff α and β are conjugate, and $\pi_0([G, H])$ is the set of conjugacy classes of homomorphisms.

Finite groupoids

- ▶ Many things about Morava E -theory are more convenient using groupoids.
- ▶ A *groupoid* is a category G in which all morphisms are invertible.
- ▶ Say G is *finite* if all Hom sets $G(a, b)$ are finite, and the set $\pi_0(G)$ of isomorphism classes is finite.
- ▶ If so, we can choose a_1, \dots, a_m containing one element of each isomorphism class, and put $G_i = G(a_i, a_i)$, and we get $BG \simeq \coprod_i BG_i$.
- ▶ Thus $E^*(BG) = \prod_i E^*(BG_i)$, which is a finitely generated E^* -module.
- ▶ Any group can be regarded as a groupoid with one object.
- ▶ A *representation* of G is a functor V from G to the category \mathcal{V} of finite-dimensional complex vector spaces.
- ▶ This again gives spaces $\text{Flag}_k(V)$ and $P(V) = \text{Flag}_1(V)$ over BG .
- ▶ Given groupoids G and H , the functor category $[G, H]$ is also a groupoid.
- ▶ If G, H are groups then $\text{obj}([G, H]) = \text{Hom}(G, H)$ and morphisms $\alpha \rightarrow \beta$ in $[G, H]$ are elements $h \in H$ with $\beta(g) = h\alpha(g)h^{-1}$ for all $g \in G$.
- ▶ So $\alpha \simeq \beta$ iff α and β are conjugate, and $\pi_0([G, H])$ is the set of conjugacy classes of homomorphisms.

- ▶ Many things about Morava E -theory are more convenient using groupoids.
- ▶ A *groupoid* is a category G in which all morphisms are invertible.
- ▶ Say G is *finite* if all Hom sets $G(a, b)$ are finite, and the set $\pi_0(G)$ of isomorphism classes is finite.
- ▶ If so, we can choose a_1, \dots, a_m containing one element of each isomorphism class, and put $G_i = G(a_i, a_i)$, and we get $BG \simeq \coprod_i BG_i$.
- ▶ Thus $E^*(BG) = \prod_i E^*(BG_i)$, which is a finitely generated E^* -module.
- ▶ Any group can be regarded as a groupoid with one object.
- ▶ A *representation* of G is a functor V from G to the category \mathcal{V} of finite-dimensional complex vector spaces.
- ▶ This again gives spaces $\text{Flag}_k(V)$ and $P(V) = \text{Flag}_1(V)$ over BG .
- ▶ Given groupoids G and H , the functor category $[G, H]$ is also a groupoid.
- ▶ If G, H are groups then $\text{obj}([G, H]) = \text{Hom}(G, H)$ and morphisms $\alpha \rightarrow \beta$ in $[G, H]$ are elements $h \in H$ with $\beta(g) = h\alpha(g)h^{-1}$ for all $g \in G$.
- ▶ So $\alpha \simeq \beta$ iff α and β are conjugate, and $\pi_0([G, H])$ is the set of conjugacy classes of homomorphisms.

- ▶ Many things about Morava E -theory are more convenient using groupoids.
- ▶ A *groupoid* is a category G in which all morphisms are invertible.
- ▶ Say G is *finite* if all Hom sets $G(a, b)$ are finite, and the set $\pi_0(G)$ of isomorphism classes is finite.
- ▶ If so, we can choose a_1, \dots, a_m containing one element of each isomorphism class, and put $G_i = G(a_i, a_i)$, and we get $BG \simeq \coprod_i BG_i$.
- ▶ Thus $E^*(BG) = \prod_i E^*(BG_i)$, which is a finitely generated E^* -module.
- ▶ Any group can be regarded as a groupoid with one object.
- ▶ A *representation* of G is a functor V from G to the category \mathcal{V} of finite-dimensional complex vector spaces.
- ▶ This again gives spaces $\text{Flag}_k(V)$ and $P(V) = \text{Flag}_1(V)$ over BG .
- ▶ Given groupoids G and H , the functor category $[G, H]$ is also a groupoid.
- ▶ If G, H are groups then $\text{obj}([G, H]) = \text{Hom}(G, H)$ and morphisms $\alpha \rightarrow \beta$ in $[G, H]$ are elements $h \in H$ with $\beta(g) = h\alpha(g)h^{-1}$ for all $g \in G$.
- ▶ So $\alpha \simeq \beta$ iff α and β are conjugate, and $\pi_0([G, H])$ is the set of conjugacy classes of homomorphisms.

Naive groupoid duality

- ▶ For a finite groupoid G put $M(G) = \mathbb{Q}\{\pi_0(G)\}$ and $M^*(G) = \text{Hom}(M(G), \mathbb{Q}) = \text{Map}(\pi_0(G), \mathbb{Q})$.
- ▶ Define an inner product on $M(G)$ by $([a], [b])_G = |G(a, b)|$ (so $([a], [b]) = 0$ unless $a \simeq b$).
- ▶ The induced inner product on $M^*(G)$ is $\langle f, g \rangle_G = \sum_{i=1}^r |G(a_i, a_i)|^{-1} f(a_i) g(a_i)$, where a_1, \dots, a_r contains one member of each isomorphism class.
- ▶ This is also $\langle f, g \rangle_G = \theta(fg)$, where $\theta(h) = \sum_i |G(a_i, a_i)|^{-1} h(a_i)$.
- ▶ Given $q: G \rightarrow H$ we define $q_!: M(G) \rightarrow M(H)$ by $q_!([a]) = [q(a)]$, and $q^*: M^*(H) \rightarrow M^*(G)$ by $q^*(g)(a) = g(q(a))$.
- ▶ Define $q^*: M(H) \rightarrow M(G)$ and $q_!: M^*(G) \rightarrow M^*(H)$ to be adjoint, so $(q_!(u), v)_H = (u, q^*(v))_G$ and $\langle q_!(f), g \rangle_H = \langle f, q^*(g) \rangle_G$.
- ▶ This is compatible with the isomorphisms $M(G) \simeq M^*(G) \simeq \text{Hom}(M(G), \mathbb{Q})$.

Naive groupoid duality

- ▶ For a finite groupoid G put $M(G) = \mathbb{Q}\{\pi_0(G)\}$ and $M^*(G) = \text{Hom}(M(G), \mathbb{Q}) = \text{Map}(\pi_0(G), \mathbb{Q})$.
- ▶ Define an inner product on $M(G)$ by $([a], [b])_G = |G(a, b)|$ (so $([a], [b]) = 0$ unless $a \simeq b$).
- ▶ The induced inner product on $M^*(G)$ is $\langle f, g \rangle_G = \sum_{i=1}^r |G(a_i, a_i)|^{-1} f(a_i) g(a_i)$, where a_1, \dots, a_r contains one member of each isomorphism class.
- ▶ This is also $\langle f, g \rangle_G = \theta(fg)$, where $\theta(h) = \sum_i |G(a_i, a_i)|^{-1} h(a_i)$.
- ▶ Given $q: G \rightarrow H$ we define $q_!: M(G) \rightarrow M(H)$ by $q_!([a]) = [q(a)]$, and $q^*: M^*(H) \rightarrow M^*(G)$ by $q^*(g)(a) = g(q(a))$.
- ▶ Define $q^*: M(H) \rightarrow M(G)$ and $q_!: M^*(G) \rightarrow M^*(H)$ to be adjoint, so $(q_!(u), v)_H = (u, q^*(v))_G$ and $(q_!(f), g)_H = \langle f, q^*(g) \rangle_G$.
- ▶ This is compatible with the isomorphisms $M(G) \simeq M^*(G) \simeq \text{Hom}(M(G), \mathbb{Q})$.

Naive groupoid duality

- ▶ For a finite groupoid G put $M(G) = \mathbb{Q}\{\pi_0(G)\}$ and $M^*(G) = \text{Hom}(M(G), \mathbb{Q}) = \text{Map}(\pi_0(G), \mathbb{Q})$.
- ▶ Define an inner product on $M(G)$ by $([a], [b])_G = |G(a, b)|$ (so $([a], [b]) = 0$ unless $a \simeq b$).
- ▶ The induced inner product on $M^*(G)$ is $\langle f, g \rangle_G = \sum_{i=1}^r |G(a_i, a_i)|^{-1} f(a_i) g(a_i)$, where a_1, \dots, a_r contains one member of each isomorphism class.
- ▶ This is also $\langle f, g \rangle_G = \theta(fg)$, where $\theta(h) = \sum_i |G(a_i, a_i)|^{-1} h(a_i)$.
- ▶ Given $q: G \rightarrow H$ we define $q_!: M(G) \rightarrow M(H)$ by $q_!([a]) = [q(a)]$, and $q^*: M^*(H) \rightarrow M^*(G)$ by $q^*(g)(a) = g(q(a))$.
- ▶ Define $q^*: M(H) \rightarrow M(G)$ and $q_!: M^*(G) \rightarrow M^*(H)$ to be adjoint, so $(q_!(u), v)_H = (u, q^*(v))_G$ and $\langle q_!(f), g \rangle_H = \langle f, q^*(g) \rangle_G$.
- ▶ This is compatible with the isomorphisms $M(G) \simeq M^*(G) \simeq \text{Hom}(M(G), \mathbb{Q})$.

Naive groupoid duality

- ▶ For a finite groupoid G put $M(G) = \mathbb{Q}\{\pi_0(G)\}$ and $M^*(G) = \text{Hom}(M(G), \mathbb{Q}) = \text{Map}(\pi_0(G), \mathbb{Q})$.
- ▶ Define an inner product on $M(G)$ by $([a], [b])_G = |G(a, b)|$ (so $([a], [b]) = 0$ unless $a \simeq b$).
- ▶ The induced inner product on $M^*(G)$ is $\langle f, g \rangle_G = \sum_{i=1}^r |G(a_i, a_i)|^{-1} f(a_i) g(a_i)$, where a_1, \dots, a_r contains one member of each isomorphism class.
- ▶ This is also $\langle f, g \rangle_G = \theta(fg)$, where $\theta(h) = \sum_i |G(a_i, a_i)|^{-1} h(a_i)$.
- ▶ Given $q: G \rightarrow H$ we define $q_!: M(G) \rightarrow M(H)$ by $q_!([a]) = [q(a)]$, and $q^*: M^*(H) \rightarrow M^*(G)$ by $q^*(g)(a) = g(q(a))$.
- ▶ Define $q^*: M(H) \rightarrow M(G)$ and $q_!: M^*(G) \rightarrow M^*(H)$ to be adjoint, so $(q_!(u), v)_H = (u, q^*(v))_G$ and $\langle q_!(f), g \rangle_H = \langle f, q^*(g) \rangle_G$.
- ▶ This is compatible with the isomorphisms $M(G) \simeq M^*(G) \simeq \text{Hom}(M(G), \mathbb{Q})$.

Naive groupoid duality

- ▶ For a finite groupoid G put $M(G) = \mathbb{Q}\{\pi_0(G)\}$ and $M^*(G) = \text{Hom}(M(G), \mathbb{Q}) = \text{Map}(\pi_0(G), \mathbb{Q})$.
- ▶ Define an inner product on $M(G)$ by $([a], [b])_G = |G(a, b)|$ (so $([a], [b]) = 0$ unless $a \simeq b$).
- ▶ The induced inner product on $M^*(G)$ is $\langle f, g \rangle_G = \sum_{i=1}^r |G(a_i, a_i)|^{-1} f(a_i) g(a_i)$, where a_1, \dots, a_r contains one member of each isomorphism class.
- ▶ This is also $\langle f, g \rangle_G = \theta(fg)$, where $\theta(h) = \sum_i |G(a_i, a_i)|^{-1} h(a_i)$.
- ▶ Given $q: G \rightarrow H$ we define $q_!: M(G) \rightarrow M(H)$ by $q_!([a]) = [q(a)]$, and $q^*: M^*(H) \rightarrow M^*(G)$ by $q^*(g)(a) = g(q(a))$.
- ▶ Define $q^*: M(H) \rightarrow M(G)$ and $q_!: M^*(G) \rightarrow M^*(H)$ to be adjoint, so $(q_!(u), v)_H = (u, q^*(v))_G$ and $\langle q_!(f), g \rangle_H = \langle f, q^*(g) \rangle_G$.
- ▶ This is compatible with the isomorphisms $M(G) \simeq M^*(G) \simeq \text{Hom}(M(G), \mathbb{Q})$.

Naive groupoid duality

- ▶ For a finite groupoid G put $M(G) = \mathbb{Q}\{\pi_0(G)\}$ and $M^*(G) = \text{Hom}(M(G), \mathbb{Q}) = \text{Map}(\pi_0(G), \mathbb{Q})$.
- ▶ Define an inner product on $M(G)$ by $([a], [b])_G = |G(a, b)|$ (so $([a], [b]) = 0$ unless $a \simeq b$).
- ▶ The induced inner product on $M^*(G)$ is $\langle f, g \rangle_G = \sum_{i=1}^r |G(a_i, a_i)|^{-1} f(a_i) g(a_i)$, where a_1, \dots, a_r contains one member of each isomorphism class.
- ▶ This is also $\langle f, g \rangle_G = \theta(fg)$, where $\theta(h) = \sum_i |G(a_i, a_i)|^{-1} h(a_i)$.
- ▶ Given $q: G \rightarrow H$ we define $q_!: M(G) \rightarrow M(H)$ by $q_!([a]) = [q(a)]$, and $q^*: M^*(H) \rightarrow M^*(G)$ by $q^*(g)(a) = g(q(a))$.
- ▶ Define $q^*: M(H) \rightarrow M(G)$ and $q_!: M^*(G) \rightarrow M^*(H)$ to be adjoint, so $(q_!(u), v)_H = (u, q^*(v))_G$ and $\langle q_!(f), g \rangle_H = \langle f, q^*(g) \rangle_G$.
- ▶ This is compatible with the isomorphisms $M(G) \simeq M^*(G) \simeq \text{Hom}(M(G), \mathbb{Q})$.

Naive groupoid duality

- ▶ For a finite groupoid G put $M(G) = \mathbb{Q}\{\pi_0(G)\}$ and $M^*(G) = \text{Hom}(M(G), \mathbb{Q}) = \text{Map}(\pi_0(G), \mathbb{Q})$.
- ▶ Define an inner product on $M(G)$ by $([a], [b])_G = |G(a, b)|$ (so $([a], [b]) = 0$ unless $a \simeq b$).
- ▶ The induced inner product on $M^*(G)$ is $\langle f, g \rangle_G = \sum_{i=1}^r |G(a_i, a_i)|^{-1} f(a_i) g(a_i)$, where a_1, \dots, a_r contains one member of each isomorphism class.
- ▶ This is also $\langle f, g \rangle_G = \theta(fg)$, where $\theta(h) = \sum_i |G(a_i, a_i)|^{-1} h(a_i)$.
- ▶ Given $q: G \rightarrow H$ we define $q_!: M(G) \rightarrow M(H)$ by $q_!([a]) = [q(a)]$, and $q^*: M^*(H) \rightarrow M^*(G)$ by $q^*(g)(a) = g(q(a))$.
- ▶ Define $q^*: M(H) \rightarrow M(G)$ and $q_!: M^*(G) \rightarrow M^*(H)$ to be adjoint, so $(q_!(u), v)_H = (u, q^*(v))_G$ and $\langle q_!(f), g \rangle_H = \langle f, q^*(g) \rangle_G$.
- ▶ This is compatible with the isomorphisms $M(G) \simeq M^*(G) \simeq \text{Hom}(M(G), \mathbb{Q})$.

Naive groupoid duality

- ▶ For a finite groupoid G put $M(G) = \mathbb{Q}\{\pi_0(G)\}$ and $M^*(G) = \text{Hom}(M(G), \mathbb{Q}) = \text{Map}(\pi_0(G), \mathbb{Q})$.
- ▶ Define an inner product on $M(G)$ by $([a], [b])_G = |G(a, b)|$ (so $([a], [b]) = 0$ unless $a \simeq b$).
- ▶ The induced inner product on $M^*(G)$ is $\langle f, g \rangle_G = \sum_{i=1}^r |G(a_i, a_i)|^{-1} f(a_i) g(a_i)$, where a_1, \dots, a_r contains one member of each isomorphism class.
- ▶ This is also $\langle f, g \rangle_G = \theta(fg)$, where $\theta(h) = \sum_i |G(a_i, a_i)|^{-1} h(a_i)$.
- ▶ Given $q: G \rightarrow H$ we define $q_!: M(G) \rightarrow M(H)$ by $q_!([a]) = [q(a)]$, and $q^*: M^*(H) \rightarrow M^*(G)$ by $q^*(g)(a) = g(q(a))$.
- ▶ Define $q^*: M(H) \rightarrow M(G)$ and $q_!: M^*(G) \rightarrow M^*(H)$ to be adjoint, so $(q_!(u), v)_H = (u, q^*(v))_G$ and $\langle q_!(f), g \rangle_H = \langle f, q^*(g) \rangle_G$.
- ▶ This is compatible with the isomorphisms $M(G) \simeq M^*(G) \simeq \text{Hom}(M(G), \mathbb{Q})$.

Generalised characters

- ▶ Fix a prime p and $n > 0$ and let E be Morava E -theory.
- ▶ Then $[p^k]_E(x) = g_k(x)h_k(x)$, where $h_k(x) \in E^0[[x]]^\times$ and $g_k(x) \in E^0[x]$ is a monic polynomial of degree p^{nk} and $E^0(BC_{p^k}) = E^0[x]/g_k(x)$.
- ▶ Construct L from $\mathbb{Q} \otimes E^0$ by adjoining a full set of roots of $g_k(x)$ for all k .
- ▶ Put $\mathbb{Z}/p^\infty = \lim_{\rightarrow k} \mathbb{Z}/p^k = \mathbb{Z}[\frac{1}{p}]/\mathbb{Z} = \mathbb{Q}/\mathbb{Z}_{(p)} = \mathbb{Q}_p/\mathbb{Z}_p = \bigcup_k \sqrt[p^k]{1} \subset S^1$.
(Exercise: $\text{Hom}(\mathbb{Z}/p^\infty, \mathbb{Z}/p^\infty) \simeq \mathbb{Z}_p \simeq \text{Hom}(\mathbb{Z}/p^\infty, S^1)$.)
- ▶ Put $\Theta = \{\text{all roots of all } g_k(x)\} \subset L$. This is a group under $+_E$, iso to $(\mathbb{Z}/p^\infty)^n$, analogous to the formal group scheme $\text{spf}(E^0(\mathbb{C}P^\infty))$.
- ▶ Put $\Theta^* = \text{Hom}(\Theta, S^1) \simeq \mathbb{Z}_p^n$, regarded as a groupoid with one object.
- ▶ Put $\Lambda G = [\Theta^*, G] = \lim_{\rightarrow k} [\Theta^*/p^k, G]$, $C(G) = L \otimes M^* \Lambda G = \text{Map}(\pi_0 \Lambda G, L)$.
- ▶ Recall $E^0(B(\Theta^*/p^k)) = E^0[[x_1, \dots, x_n]]/(g_k(x_1), \dots, g_k(x_n))$; there is a canonical map ϕ_k from this to L .
- ▶ Thus any $u: \Theta^*/p^k \rightarrow G$ gives $\phi_k \circ E^0(Bu): E^0 BG \rightarrow L$. Assembling these gives $\chi: L \otimes_{E^0} E^0(BG) \rightarrow C(G)$.
- ▶ Theorem (Hopkins, Kuhn, Ravenel): χ is an isomorphism.

Generalised characters

- ▶ Fix a prime p and $n > 0$ and let E be Morava E -theory.
- ▶ Then $[p^k]_E(x) = g_k(x)h_k(x)$, where $h_k(x) \in E^0[[x]]^\times$ and $g_k(x) \in E^0[x]$ is a monic polynomial of degree p^{nk} and $E^0(BC_{p^k}) = E^0[x]/g_k(x)$.
- ▶ Construct L from $\mathbb{Q} \otimes E^0$ by adjoining a full set of roots of $g_k(x)$ for all k .
- ▶ Put $\mathbb{Z}/p^\infty = \lim_{\rightarrow k} \mathbb{Z}/p^k = \mathbb{Z}[\frac{1}{p}]/\mathbb{Z} = \mathbb{Q}/\mathbb{Z}_{(p)} = \mathbb{Q}_p/\mathbb{Z}_p = \bigcup_k \sqrt[p^k]{1} \subset S^1$.
(Exercise: $\text{Hom}(\mathbb{Z}/p^\infty, \mathbb{Z}/p^\infty) \simeq \mathbb{Z}_p \simeq \text{Hom}(\mathbb{Z}/p^\infty, S^1)$.)
- ▶ Put $\Theta = \{\text{all roots of all } g_k(x)\} \subset L$. This is a group under $+_E$, iso to $(\mathbb{Z}/p^\infty)^n$, analogous to the formal group scheme $\text{spf}(E^0(\mathbb{C}P^\infty))$.
- ▶ Put $\Theta^* = \text{Hom}(\Theta, S^1) \simeq \mathbb{Z}_p^n$, regarded as a groupoid with one object.
- ▶ Put $\Lambda G = [\Theta^*, G] = \lim_{\rightarrow k} [\Theta^*/p^k, G]$, $C(G) = L \otimes M^* \Lambda G = \text{Map}(\pi_0 \Lambda G, L)$.
- ▶ Recall $E^0(B(\Theta^*/p^k)) = E^0[[x_1, \dots, x_n]]/(g_k(x_1), \dots, g_k(x_n))$; there is a canonical map ϕ_k from this to L .
- ▶ Thus any $u: \Theta^*/p^k \rightarrow G$ gives $\phi_k \circ E^0(Bu): E^0BG \rightarrow L$. Assembling these gives $\chi: L \otimes_{E^0} E^0(BG) \rightarrow C(G)$.
- ▶ Theorem (Hopkins, Kuhn, Ravenel): χ is an isomorphism.

Generalised characters

- ▶ Fix a prime p and $n > 0$ and let E be Morava E -theory.
- ▶ Then $[p^k]_E(x) = g_k(x)h_k(x)$, where $h_k(x) \in E^0[[x]]^\times$ and $g_k(x) \in E^0[x]$ is a monic polynomial of degree p^{nk} and $E^0(BC_{p^k}) = E^0[x]/g_k(x)$.
- ▶ Construct L from $\mathbb{Q} \otimes E^0$ by adjoining a full set of roots of $g_k(x)$ for all k .
- ▶ Put $\mathbb{Z}/p^\infty = \lim_{\rightarrow k} \mathbb{Z}/p^k = \mathbb{Z}[\frac{1}{p}]/\mathbb{Z} = \mathbb{Q}/\mathbb{Z}_{(p)} = \mathbb{Q}_p/\mathbb{Z}_p = \bigcup_k \sqrt[p^k]{1} \subset S^1$.
(Exercise: $\text{Hom}(\mathbb{Z}/p^\infty, \mathbb{Z}/p^\infty) \simeq \mathbb{Z}_p \simeq \text{Hom}(\mathbb{Z}/p^\infty, S^1)$.)
- ▶ Put $\Theta = \{\text{all roots of all } g_k(x)\} \subset L$. This is a group under $+_E$, iso to $(\mathbb{Z}/p^\infty)^n$, analogous to the formal group scheme $\text{spf}(E^0(\mathbb{C}P^\infty))$.
- ▶ Put $\Theta^* = \text{Hom}(\Theta, S^1) \simeq \mathbb{Z}_p^n$, regarded as a groupoid with one object.
- ▶ Put $\Lambda G = [\Theta^*, G] = \lim_{\rightarrow k} [\Theta^*/p^k, G]$, $C(G) = L \otimes M^* \Lambda G = \text{Map}(\pi_0 \Lambda G, L)$.
- ▶ Recall $E^0(B(\Theta^*/p^k)) = E^0[[x_1, \dots, x_n]]/(g_k(x_1), \dots, g_k(x_n))$; there is a canonical map ϕ_k from this to L .
- ▶ Thus any $u: \Theta^*/p^k \rightarrow G$ gives $\phi_k \circ E^0(Bu): E^0 BG \rightarrow L$. Assembling these gives $\chi: L \otimes_{E^0} E^0(BG) \rightarrow C(G)$.
- ▶ Theorem (Hopkins, Kuhn, Ravenel): χ is an isomorphism.

Generalised characters

- ▶ Fix a prime p and $n > 0$ and let E be Morava E -theory.
- ▶ Then $[p^k]_E(x) = g_k(x)h_k(x)$, where $h_k(x) \in E^0[[x]]^\times$ and $g_k(x) \in E^0[x]$ is a monic polynomial of degree p^{nk} and $E^0(BC_{p^k}) = E^0[x]/g_k(x)$.
- ▶ Construct L from $\mathbb{Q} \otimes E^0$ by adjoining a full set of roots of $g_k(x)$ for all k .
- ▶ Put $\mathbb{Z}/p^\infty = \lim_{\rightarrow k} \mathbb{Z}/p^k = \mathbb{Z}[\frac{1}{p}]/\mathbb{Z} = \mathbb{Q}/\mathbb{Z}_{(p)} = \mathbb{Q}_p/\mathbb{Z}_p = \bigcup_k \sqrt[p^k]{1} \subset S^1$.
(Exercise: $\text{Hom}(\mathbb{Z}/p^\infty, \mathbb{Z}/p^\infty) \simeq \mathbb{Z}_p \simeq \text{Hom}(\mathbb{Z}/p^\infty, S^1)$.)
- ▶ Put $\Theta = \{\text{all roots of all } g_k(x)\} \subset L$. This is a group under $+_E$, iso to $(\mathbb{Z}/p^\infty)^n$, analogous to the formal group scheme $\text{spf}(E^0(\mathbb{C}P^\infty))$.
- ▶ Put $\Theta^* = \text{Hom}(\Theta, S^1) \simeq \mathbb{Z}_p^n$, regarded as a groupoid with one object.
- ▶ Put $\Lambda G = [\Theta^*, G] = \lim_{\rightarrow k} [\Theta^*/p^k, G]$, $C(G) = L \otimes M^* \Lambda G = \text{Map}(\pi_0 \Lambda G, L)$.
- ▶ Recall $E^0(B(\Theta^*/p^k)) = E^0[[x_1, \dots, x_n]]/(g_k(x_1), \dots, g_k(x_n))$; there is a canonical map ϕ_k from this to L .
- ▶ Thus any $u: \Theta^*/p^k \rightarrow G$ gives $\phi_k \circ E^0(Bu): E^0BG \rightarrow L$.
Assembling these gives $\chi: L \otimes_{E^0} E^0(BG) \rightarrow C(G)$.
- ▶ Theorem (Hopkins, Kuhn, Ravenel): χ is an isomorphism.

Generalised characters

- ▶ Fix a prime p and $n > 0$ and let E be Morava E -theory.
- ▶ Then $[p^k]_E(x) = g_k(x)h_k(x)$, where $h_k(x) \in E^0[[x]]^\times$ and $g_k(x) \in E^0[x]$ is a monic polynomial of degree p^{nk} and $E^0(BC_{p^k}) = E^0[x]/g_k(x)$.
- ▶ Construct L from $\mathbb{Q} \otimes E^0$ by adjoining a full set of roots of $g_k(x)$ for all k .
- ▶ Put $\mathbb{Z}/p^\infty = \lim_{\rightarrow k} \mathbb{Z}/p^k = \mathbb{Z}[\frac{1}{p}]/\mathbb{Z} = \mathbb{Q}/\mathbb{Z}_{(p)} = \mathbb{Q}_p/\mathbb{Z}_p = \bigcup_k \sqrt[p^k]{1} \subset S^1$.
(Exercise: $\text{Hom}(\mathbb{Z}/p^\infty, \mathbb{Z}/p^\infty) \simeq \mathbb{Z}_p \simeq \text{Hom}(\mathbb{Z}/p^\infty, S^1)$.)
- ▶ Put $\Theta = \{\text{all roots of all } g_k(x)\} \subset L$. This is a group under $+_E$, iso to $(\mathbb{Z}/p^\infty)^n$, analogous to the formal group scheme $\text{spf}(E^0(\mathbb{C}P^\infty))$.
- ▶ Put $\Theta^* = \text{Hom}(\Theta, S^1) \simeq \mathbb{Z}_p^n$, regarded as a groupoid with one object.
- ▶ Put $\Lambda G = [\Theta^*, G] = \lim_{\rightarrow k} [\Theta^*/p^k, G]$, $C(G) = L \otimes M^* \Lambda G = \text{Map}(\pi_0 \Lambda G, L)$.
- ▶ Recall $E^0(B(\Theta^*/p^k)) = E^0[[x_1, \dots, x_n]]/(g_k(x_1), \dots, g_k(x_n))$; there is a canonical map ϕ_k from this to L .
- ▶ Thus any $u: \Theta^*/p^k \rightarrow G$ gives $\phi_k \circ E^0(Bu): E^0BG \rightarrow L$. Assembling these gives $\chi: L \otimes_{E^0} E^0(BG) \rightarrow C(G)$.
- ▶ Theorem (Hopkins, Kuhn, Ravenel): χ is an isomorphism.

Generalised characters

- ▶ Fix a prime p and $n > 0$ and let E be Morava E -theory.
- ▶ Then $[p^k]_E(x) = g_k(x)h_k(x)$, where $h_k(x) \in E^0[[x]]^\times$ and $g_k(x) \in E^0[x]$ is a monic polynomial of degree p^{nk} and $E^0(BC_{p^k}) = E^0[x]/g_k(x)$.
- ▶ Construct L from $\mathbb{Q} \otimes E^0$ by adjoining a full set of roots of $g_k(x)$ for all k .
- ▶ Put $\mathbb{Z}/p^\infty = \lim_{\rightarrow k} \mathbb{Z}/p^k = \mathbb{Z}[\frac{1}{p}]/\mathbb{Z} = \mathbb{Q}/\mathbb{Z}_{(p)} = \mathbb{Q}_p/\mathbb{Z}_p = \bigcup_k \sqrt[p^k]{1} \subset S^1$.
(Exercise: $\text{Hom}(\mathbb{Z}/p^\infty, \mathbb{Z}/p^\infty) \simeq \mathbb{Z}_p \simeq \text{Hom}(\mathbb{Z}/p^\infty, S^1)$.)
- ▶ Put $\Theta = \{\text{all roots of all } g_k(x)\} \subset L$. This is a group under $+_E$, iso to $(\mathbb{Z}/p^\infty)^n$, analogous to the formal group scheme $\text{spf}(E^0(\mathbb{C}P^\infty))$.
- ▶ Put $\Theta^* = \text{Hom}(\Theta, S^1) \simeq \mathbb{Z}_p^n$, regarded as a groupoid with one object.
- ▶ Put $\Lambda G = [\Theta^*, G] = \lim_{\rightarrow k} [\Theta^*/p^k, G]$, $C(G) = L \otimes M^* \Lambda G = \text{Map}(\pi_0 \Lambda G, L)$.
- ▶ Recall $E^0(B(\Theta^*/p^k)) = E^0[[x_1, \dots, x_n]]/(g_k(x_1), \dots, g_k(x_n))$; there is a canonical map ϕ_k from this to L .
- ▶ Thus any $u: \Theta^*/p^k \rightarrow G$ gives $\phi_k \circ E^0(Bu): E^0BG \rightarrow L$. Assembling these gives $\chi: L \otimes_{E^0} E^0(BG) \rightarrow C(G)$.
- ▶ Theorem (Hopkins, Kuhn, Ravenel): χ is an isomorphism.

Generalised characters

- ▶ Fix a prime p and $n > 0$ and let E be Morava E -theory.
- ▶ Then $[p^k]_E(x) = g_k(x)h_k(x)$, where $h_k(x) \in E^0[[x]]^\times$ and $g_k(x) \in E^0[x]$ is a monic polynomial of degree p^{nk} and $E^0(BC_{p^k}) = E^0[x]/g_k(x)$.
- ▶ Construct L from $\mathbb{Q} \otimes E^0$ by adjoining a full set of roots of $g_k(x)$ for all k .
- ▶ Put $\mathbb{Z}/p^\infty = \lim_{\rightarrow k} \mathbb{Z}/p^k = \mathbb{Z}[\frac{1}{p}]/\mathbb{Z} = \mathbb{Q}/\mathbb{Z}_{(p)} = \mathbb{Q}_p/\mathbb{Z}_p = \bigcup_k \sqrt[p^k]{1} \subset S^1$.
(Exercise: $\text{Hom}(\mathbb{Z}/p^\infty, \mathbb{Z}/p^\infty) \simeq \mathbb{Z}_p \simeq \text{Hom}(\mathbb{Z}/p^\infty, S^1)$.)
- ▶ Put $\Theta = \{\text{all roots of all } g_k(x)\} \subset L$. This is a group under $+_E$, iso to $(\mathbb{Z}/p^\infty)^n$, analogous to the formal group scheme $\text{spf}(E^0(\mathbb{C}P^\infty))$.
- ▶ Put $\Theta^* = \text{Hom}(\Theta, S^1) \simeq \mathbb{Z}_p^n$, regarded as a groupoid with one object.
- ▶ Put $\Lambda G = [\Theta^*, G] = \lim_{\rightarrow k} [\Theta^*/p^k, G]$, $C(G) = L \otimes M^* \Lambda G = \text{Map}(\pi_0 \Lambda G, L)$.
- ▶ Recall $E^0(B(\Theta^*/p^k)) = E^0[[x_1, \dots, x_n]]/(g_k(x_1), \dots, g_k(x_n))$; there is a canonical map ϕ_k from this to L .
- ▶ Thus any $u: \Theta^*/p^k \rightarrow G$ gives $\phi_k \circ E^0(Bu): E^0BG \rightarrow L$.
Assembling these gives $\chi: L \otimes_{E^0} E^0(BG) \rightarrow C(G)$.
- ▶ Theorem (Hopkins, Kuhn, Ravenel): χ is an isomorphism.

Generalised characters

- ▶ Fix a prime p and $n > 0$ and let E be Morava E -theory.
- ▶ Then $[p^k]_E(x) = g_k(x)h_k(x)$, where $h_k(x) \in E^0[[x]]^\times$ and $g_k(x) \in E^0[x]$ is a monic polynomial of degree p^{nk} and $E^0(BC_{p^k}) = E^0[x]/g_k(x)$.
- ▶ Construct L from $\mathbb{Q} \otimes E^0$ by adjoining a full set of roots of $g_k(x)$ for all k .
- ▶ Put $\mathbb{Z}/p^\infty = \lim_{\rightarrow k} \mathbb{Z}/p^k = \mathbb{Z}[\frac{1}{p}]/\mathbb{Z} = \mathbb{Q}/\mathbb{Z}_{(p)} = \mathbb{Q}_p/\mathbb{Z}_p = \bigcup_k \sqrt[p^k]{1} \subset S^1$.
(Exercise: $\text{Hom}(\mathbb{Z}/p^\infty, \mathbb{Z}/p^\infty) \simeq \mathbb{Z}_p \simeq \text{Hom}(\mathbb{Z}/p^\infty, S^1)$.)
- ▶ Put $\Theta = \{\text{all roots of all } g_k(x)\} \subset L$. This is a group under $+_E$, iso to $(\mathbb{Z}/p^\infty)^n$, analogous to the formal group scheme $\text{spf}(E^0(\mathbb{C}P^\infty))$.
- ▶ Put $\Theta^* = \text{Hom}(\Theta, S^1) \simeq \mathbb{Z}_p^n$, regarded as a groupoid with one object.
- ▶ Put $\Lambda G = [\Theta^*, G] = \lim_{\rightarrow k} [\Theta^*/p^k, G]$, $C(G) = L \otimes M^* \Lambda G = \text{Map}(\pi_0 \Lambda G, L)$.
- ▶ Recall $E^0(B(\Theta^*/p^k)) = E^0[[x_1, \dots, x_n]]/(g_k(x_1), \dots, g_k(x_n))$; there is a canonical map ϕ_k from this to L .
- ▶ Thus any $u: \Theta^*/p^k \rightarrow G$ gives $\phi_k \circ E^0(Bu): E^0BG \rightarrow L$.
Assembling these gives $\chi: L \otimes_{E^0} E^0(BG) \rightarrow C(G)$.
- ▶ Theorem (Hopkins, Kuhn, Ravenel): χ is an isomorphism.

Generalised characters

- ▶ Fix a prime p and $n > 0$ and let E be Morava E -theory.
- ▶ Then $[p^k]_E(x) = g_k(x)h_k(x)$, where $h_k(x) \in E^0[[x]]^\times$ and $g_k(x) \in E^0[x]$ is a monic polynomial of degree p^{nk} and $E^0(BC_{p^k}) = E^0[x]/g_k(x)$.
- ▶ Construct L from $\mathbb{Q} \otimes E^0$ by adjoining a full set of roots of $g_k(x)$ for all k .
- ▶ Put $\mathbb{Z}/p^\infty = \lim_{\rightarrow k} \mathbb{Z}/p^k = \mathbb{Z}[\frac{1}{p}]/\mathbb{Z} = \mathbb{Q}/\mathbb{Z}_{(p)} = \mathbb{Q}_p/\mathbb{Z}_p = \bigcup_k \sqrt[p^k]{1} \subset S^1$.
(Exercise: $\text{Hom}(\mathbb{Z}/p^\infty, \mathbb{Z}/p^\infty) \simeq \mathbb{Z}_p \simeq \text{Hom}(\mathbb{Z}/p^\infty, S^1)$.)
- ▶ Put $\Theta = \{\text{all roots of all } g_k(x)\} \subset L$. This is a group under $+_E$, iso to $(\mathbb{Z}/p^\infty)^n$, analogous to the formal group scheme $\text{spf}(E^0(\mathbb{C}P^\infty))$.
- ▶ Put $\Theta^* = \text{Hom}(\Theta, S^1) \simeq \mathbb{Z}_p^n$, regarded as a groupoid with one object.
- ▶ Put $\Lambda G = [\Theta^*, G] = \lim_{\rightarrow k} [\Theta^*/p^k, G]$, $C(G) = L \otimes M^* \Lambda G = \text{Map}(\pi_0 \Lambda G, L)$.
- ▶ Recall $E^0(B(\Theta^*/p^k)) = E^0[[x_1, \dots, x_n]]/(g_k(x_1), \dots, g_k(x_n))$; there is a canonical map ϕ_k from this to L .
- ▶ Thus any $u: \Theta^*/p^k \rightarrow G$ gives $\phi_k \circ E^0(Bu): E^0BG \rightarrow L$.
Assembling these gives $\chi: L \otimes_{E^0} E^0(BG) \rightarrow C(G)$.
- ▶ Theorem (Hopkins, Kuhn, Ravenel): χ is an isomorphism.

Generalised characters

- ▶ Fix a prime p and $n > 0$ and let E be Morava E -theory.
- ▶ Then $[p^k]_E(x) = g_k(x)h_k(x)$, where $h_k(x) \in E^0[[x]]^\times$ and $g_k(x) \in E^0[x]$ is a monic polynomial of degree p^{nk} and $E^0(BC_{p^k}) = E^0[x]/g_k(x)$.
- ▶ Construct L from $\mathbb{Q} \otimes E^0$ by adjoining a full set of roots of $g_k(x)$ for all k .
- ▶ Put $\mathbb{Z}/p^\infty = \lim_{\rightarrow k} \mathbb{Z}/p^k = \mathbb{Z}[\frac{1}{p}]/\mathbb{Z} = \mathbb{Q}/\mathbb{Z}_{(p)} = \mathbb{Q}_p/\mathbb{Z}_p = \bigcup_k \sqrt[p^k]{1} \subset S^1$.
(Exercise: $\text{Hom}(\mathbb{Z}/p^\infty, \mathbb{Z}/p^\infty) \simeq \mathbb{Z}_p \simeq \text{Hom}(\mathbb{Z}/p^\infty, S^1)$.)
- ▶ Put $\Theta = \{\text{all roots of all } g_k(x)\} \subset L$. This is a group under $+_E$, iso to $(\mathbb{Z}/p^\infty)^n$, analogous to the formal group scheme $\text{spf}(E^0(\mathbb{C}P^\infty))$.
- ▶ Put $\Theta^* = \text{Hom}(\Theta, S^1) \simeq \mathbb{Z}_p^n$, regarded as a groupoid with one object.
- ▶ Put $\Lambda G = [\Theta^*, G] = \lim_{\rightarrow k} [\Theta^*/p^k, G]$, $C(G) = L \otimes M^* \Lambda G = \text{Map}(\pi_0 \Lambda G, L)$.
- ▶ Recall $E^0(B(\Theta^*/p^k)) = E^0[[x_1, \dots, x_n]]/(g_k(x_1), \dots, g_k(x_n))$; there is a canonical map ϕ_k from this to L .
- ▶ Thus any $u: \Theta^*/p^k \rightarrow G$ gives $\phi_k \circ E^0(Bu): E^0BG \rightarrow L$. Assembling these gives $\chi: L \otimes_{E^0} E^0(BG) \rightarrow C(G)$.
- ▶ Theorem (Hopkins, Kuhn, Ravenel): χ is an isomorphism.

Generalised characters

- ▶ Fix a prime p and $n > 0$ and let E be Morava E -theory.
- ▶ Then $[p^k]_E(x) = g_k(x)h_k(x)$, where $h_k(x) \in E^0[[x]]^\times$ and $g_k(x) \in E^0[x]$ is a monic polynomial of degree p^{nk} and $E^0(BC_{p^k}) = E^0[x]/g_k(x)$.
- ▶ Construct L from $\mathbb{Q} \otimes E^0$ by adjoining a full set of roots of $g_k(x)$ for all k .
- ▶ Put $\mathbb{Z}/p^\infty = \lim_{\rightarrow k} \mathbb{Z}/p^k = \mathbb{Z}[\frac{1}{p}]/\mathbb{Z} = \mathbb{Q}/\mathbb{Z}_{(p)} = \mathbb{Q}_p/\mathbb{Z}_p = \bigcup_k \sqrt[p^k]{1} \subset S^1$.
(Exercise: $\text{Hom}(\mathbb{Z}/p^\infty, \mathbb{Z}/p^\infty) \simeq \mathbb{Z}_p \simeq \text{Hom}(\mathbb{Z}/p^\infty, S^1)$.)
- ▶ Put $\Theta = \{\text{all roots of all } g_k(x)\} \subset L$. This is a group under $+_E$, iso to $(\mathbb{Z}/p^\infty)^n$, analogous to the formal group scheme $\text{spf}(E^0(\mathbb{C}P^\infty))$.
- ▶ Put $\Theta^* = \text{Hom}(\Theta, S^1) \simeq \mathbb{Z}_p^n$, regarded as a groupoid with one object.
- ▶ Put $\Lambda G = [\Theta^*, G] = \lim_{\rightarrow k} [\Theta^*/p^k, G]$, $C(G) = L \otimes M^* \Lambda G = \text{Map}(\pi_0 \Lambda G, L)$.
- ▶ Recall $E^0(B(\Theta^*/p^k)) = E^0[[x_1, \dots, x_n]]/(g_k(x_1), \dots, g_k(x_n))$; there is a canonical map ϕ_k from this to L .
- ▶ Thus any $u: \Theta^*/p^k \rightarrow G$ gives $\phi_k \circ E^0(Bu): E^0 BG \rightarrow L$.
Assembling these gives $\chi: L \otimes_{E^0} E^0(BG) \rightarrow C(G)$.
- ▶ Theorem (Hopkins, Kuhn, Ravenel): χ is an isomorphism.

Generalised characters

- ▶ Fix a prime p and $n > 0$ and let E be Morava E -theory.
- ▶ Then $[p^k]_E(x) = g_k(x)h_k(x)$, where $h_k(x) \in E^0[[x]]^\times$ and $g_k(x) \in E^0[x]$ is a monic polynomial of degree p^{nk} and $E^0(BC_{p^k}) = E^0[x]/g_k(x)$.
- ▶ Construct L from $\mathbb{Q} \otimes E^0$ by adjoining a full set of roots of $g_k(x)$ for all k .
- ▶ Put $\mathbb{Z}/p^\infty = \lim_{\rightarrow k} \mathbb{Z}/p^k = \mathbb{Z}[\frac{1}{p}]/\mathbb{Z} = \mathbb{Q}/\mathbb{Z}_{(p)} = \mathbb{Q}_p/\mathbb{Z}_p = \bigcup_k \sqrt[p^k]{1} \subset S^1$.
(Exercise: $\text{Hom}(\mathbb{Z}/p^\infty, \mathbb{Z}/p^\infty) \simeq \mathbb{Z}_p \simeq \text{Hom}(\mathbb{Z}/p^\infty, S^1)$.)
- ▶ Put $\Theta = \{\text{all roots of all } g_k(x)\} \subset L$. This is a group under $+_E$, iso to $(\mathbb{Z}/p^\infty)^n$, analogous to the formal group scheme $\text{spf}(E^0(\mathbb{C}P^\infty))$.
- ▶ Put $\Theta^* = \text{Hom}(\Theta, S^1) \simeq \mathbb{Z}_p^n$, regarded as a groupoid with one object.
- ▶ Put $\Lambda G = [\Theta^*, G] = \lim_{\rightarrow k} [\Theta^*/p^k, G]$, $C(G) = L \otimes M^* \Lambda G = \text{Map}(\pi_0 \Lambda G, L)$.
- ▶ Recall $E^0(B(\Theta^*/p^k)) = E^0[[x_1, \dots, x_n]]/(g_k(x_1), \dots, g_k(x_n))$; there is a canonical map ϕ_k from this to L .
- ▶ Thus any $u: \Theta^*/p^k \rightarrow G$ gives $\phi_k \circ E^0(Bu): E^0BG \rightarrow L$.
Assembling these gives $\chi: L \otimes_{E^0} E^0(BG) \rightarrow C(G)$.
- ▶ Theorem (Hopkins, Kuhn, Ravenel): χ is an isomorphism.

Generalised characters

- ▶ Fix a prime p and $n > 0$ and let E be Morava E -theory.
- ▶ Then $[p^k]_E(x) = g_k(x)h_k(x)$, where $h_k(x) \in E^0[[x]]^\times$ and $g_k(x) \in E^0[x]$ is a monic polynomial of degree p^{nk} and $E^0(BC_{p^k}) = E^0[x]/g_k(x)$.
- ▶ Construct L from $\mathbb{Q} \otimes E^0$ by adjoining a full set of roots of $g_k(x)$ for all k .
- ▶ Put $\mathbb{Z}/p^\infty = \lim_{\rightarrow k} \mathbb{Z}/p^k = \mathbb{Z}[\frac{1}{p}]/\mathbb{Z} = \mathbb{Q}/\mathbb{Z}_{(p)} = \mathbb{Q}_p/\mathbb{Z}_p = \bigcup_k \sqrt[p^k]{1} \subset S^1$.
(Exercise: $\text{Hom}(\mathbb{Z}/p^\infty, \mathbb{Z}/p^\infty) \simeq \mathbb{Z}_p \simeq \text{Hom}(\mathbb{Z}/p^\infty, S^1)$.)
- ▶ Put $\Theta = \{\text{all roots of all } g_k(x)\} \subset L$. This is a group under $+_E$, iso to $(\mathbb{Z}/p^\infty)^n$, analogous to the formal group scheme $\text{spf}(E^0(\mathbb{C}P^\infty))$.
- ▶ Put $\Theta^* = \text{Hom}(\Theta, S^1) \simeq \mathbb{Z}_p^n$, regarded as a groupoid with one object.
- ▶ Put $\Lambda G = [\Theta^*, G] = \lim_{\rightarrow k} [\Theta^*/p^k, G]$, $C(G) = L \otimes M^* \Lambda G = \text{Map}(\pi_0 \Lambda G, L)$.
- ▶ Recall $E^0(B(\Theta^*/p^k)) = E^0[[x_1, \dots, x_n]]/(g_k(x_1), \dots, g_k(x_n))$; there is a canonical map ϕ_k from this to L .
- ▶ Thus any $u: \Theta^*/p^k \rightarrow G$ gives $\phi_k \circ E^0(Bu): E^0 BG \rightarrow L$. Assembling these gives $\chi: L \otimes_{E^0} E^0(BG) \rightarrow C(G)$.
- ▶ Theorem (Hopkins, Kuhn, Ravenel): χ is an isomorphism.

Proof of the generalised character theorem

- ▶ Reduce to the case of a finite group G .
- ▶ Generalise: for a finite G -CW complex Z , we have

$$\chi_{G,Z}: L \otimes_{E^0} E^*(Z_{hG}) \rightarrow L \otimes_{\mathbb{Q}} \left(\prod_{\theta: \Theta^* \rightarrow G} H^*(Z^{\text{image}(\theta)}; \mathbb{Q}) \right)^G$$

- ▶ Prove by calculation that $\theta_{G,Z}$ is iso when $Z = G/A$ with $A \leq G$ abelian. (Here $Z_{hG} = BA$, and $Z^{\text{image}(\theta)}$ is Z (if $\text{image}(\theta) \leq A$) or \emptyset (otherwise).)
- ▶ Deduce by Mayer-Vietoris that $\chi_{G,Z}$ is iso if Z has abelian isotropy.
- ▶ Let $F = \{\underline{W} \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$ be the space of complete flags in $\mathbb{C}[G]$, so $Z \times F$ and $Z \times F^2$ have abelian isotropy, and we have projections $p: (Z \times F)_{hG} \rightarrow Z_{hG}$ and $q_0, q_1: (Z \times F^2)_{hG} \rightarrow (Z \times F)_{hG}$
- ▶ We saw before that $E^*((Z \times F)_{hG})$ has a canonical basis $e_1 = 1, e_2, \dots, e_{n!}$ over $E^*(Z_{hG})$. Similarly, the elements $q_0^*(e_i)q_1^*(e_j)$ form a basis for $E^*((Z \times F^2)_{hG})$ over $E^*(Z_{hG})$.
- ▶ It follows that the diagram $E^*(Z_{hG}) \rightarrow E^*((Z \times F)_{hG}) \rightrightarrows E^*((Z \times F^2)_{hG})$ is an equaliser.
- ▶ Deduce the general case from this.

Proof of the generalised character theorem

- ▶ Reduce to the case of a finite group G .
- ▶ Generalise: for a finite G -CW complex Z , we have

$$\chi_{G,Z}: L \otimes_{E^0} E^*(Z_{hG}) \rightarrow L \otimes_{\mathbb{Q}} \left(\prod_{\theta: \Theta^* \rightarrow G} H^*(Z^{\text{image}(\theta)}; \mathbb{Q}) \right)^G$$

- ▶ Prove by calculation that $\theta_{G,Z}$ is iso when $Z = G/A$ with $A \leq G$ abelian. (Here $Z_{hG} = BA$, and $Z^{\text{image}(\theta)}$ is Z (if $\text{image}(\theta) \leq A$) or \emptyset (otherwise).)
- ▶ Deduce by Mayer-Vietoris that $\chi_{G,Z}$ is iso if Z has abelian isotropy.
- ▶ Let $F = \{\underline{W} \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$ be the space of complete flags in $\mathbb{C}[G]$, so $Z \times F$ and $Z \times F^2$ have abelian isotropy, and we have projections $\rho: (Z \times F)_{hG} \rightarrow Z_{hG}$ and $q_0, q_1: (Z \times F^2)_{hG} \rightarrow (Z \times F)_{hG}$
- ▶ We saw before that $E^*((Z \times F)_{hG})$ has a canonical basis $e_1 = 1, e_2, \dots, e_{n!}$ over $E^*(Z_{hG})$. Similarly, the elements $q_0^*(e_i)q_1^*(e_j)$ form a basis for $E^*((Z \times F^2)_{hG})$ over $E^*(Z_{hG})$.
- ▶ It follows that the diagram $E^*(Z_{hG}) \rightarrow E^*((Z \times F)_{hG}) \rightrightarrows E^*((Z \times F^2)_{hG})$ is an equaliser.
- ▶ Deduce the general case from this.

Proof of the generalised character theorem

- ▶ Reduce to the case of a finite group G .
- ▶ Generalise: for a finite G -CW complex Z , we have

$$\chi_{G,Z}: L \otimes_{E^0} E^*(Z_{hG}) \rightarrow L \otimes_{\mathbb{Q}} \left(\prod_{\theta: \Theta^* \rightarrow G} H^*(Z^{\text{image}(\theta)}; \mathbb{Q}) \right)^G$$

- ▶ Prove by calculation that $\theta_{G,Z}$ is iso when $Z = G/A$ with $A \leq G$ abelian. (Here $Z_{hG} = BA$, and $Z^{\text{image}(\theta)}$ is Z (if $\text{image}(\theta) \leq A$) or \emptyset (otherwise).)
- ▶ Deduce by Mayer-Vietoris that $\chi_{G,Z}$ is iso if Z has abelian isotropy.
- ▶ Let $F = \{\underline{W} \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$ be the space of complete flags in $\mathbb{C}[G]$, so $Z \times F$ and $Z \times F^2$ have abelian isotropy, and we have projections $p: (Z \times F)_{hG} \rightarrow Z_{hG}$ and $q_0, q_1: (Z \times F^2)_{hG} \rightarrow (Z \times F)_{hG}$
- ▶ We saw before that $E^*((Z \times F)_{hG})$ has a canonical basis $e_1 = 1, e_2, \dots, e_{n!}$ over $E^*(Z_{hG})$. Similarly, the elements $q_0^*(e_i)q_1^*(e_j)$ form a basis for $E^*((Z \times F^2)_{hG})$ over $E^*(Z_{hG})$.
- ▶ It follows that the diagram $E^*(Z_{hG}) \rightarrow E^*((Z \times F)_{hG}) \rightrightarrows E^*((Z \times F^2)_{hG})$ is an equaliser.
- ▶ Deduce the general case from this.

Proof of the generalised character theorem

- ▶ Reduce to the case of a finite group G .
- ▶ Generalise: for a finite G -CW complex Z , we have

$$\chi_{G,Z}: L \otimes_{E^0} E^*(Z_{hG}) \rightarrow L \otimes_{\mathbb{Q}} \left(\prod_{\theta: \Theta^* \rightarrow G} H^*(Z^{\text{image}(\theta)}; \mathbb{Q}) \right)^G$$

- ▶ Prove by calculation that $\theta_{G,Z}$ is iso when $Z = G/A$ with $A \leq G$ abelian. (Here $Z_{hG} = BA$, and $Z^{\text{image}(\theta)}$ is Z (if $\text{image}(\theta) \leq A$) or \emptyset (otherwise).)
- ▶ Deduce by Mayer-Vietoris that $\chi_{G,Z}$ is iso if Z has abelian isotropy.
- ▶ Let $F = \{\underline{W} \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$ be the space of complete flags in $\mathbb{C}[G]$, so $Z \times F$ and $Z \times F^2$ have abelian isotropy, and we have projections $p: (Z \times F)_{hG} \rightarrow Z_{hG}$ and $q_0, q_1: (Z \times F^2)_{hG} \rightarrow (Z \times F)_{hG}$
- ▶ We saw before that $E^*((Z \times F)_{hG})$ has a canonical basis $e_1 = 1, e_2, \dots, e_{n!}$ over $E^*(Z_{hG})$. Similarly, the elements $q_0^*(e_i)q_1^*(e_j)$ form a basis for $E^*((Z \times F^2)_{hG})$ over $E^*(Z_{hG})$.
- ▶ It follows that the diagram $E^*(Z_{hG}) \rightarrow E^*((Z \times F)_{hG}) \rightrightarrows E^*((Z \times F^2)_{hG})$ is an equaliser.
- ▶ Deduce the general case from this.

Proof of the generalised character theorem

- ▶ Reduce to the case of a finite group G .
- ▶ Generalise: for a finite G -CW complex Z , we have

$$\chi_{G,Z}: L \otimes_{E^0} E^*(Z_{hG}) \rightarrow L \otimes_{\mathbb{Q}} \left(\prod_{\theta: \Theta^* \rightarrow G} H^*(Z^{\text{image}(\theta)}; \mathbb{Q}) \right)^G$$

- ▶ Prove by calculation that $\theta_{G,Z}$ is iso when $Z = G/A$ with $A \leq G$ abelian. (Here $Z_{hG} = BA$, and $Z^{\text{image}(\theta)}$ is Z (if $\text{image}(\theta) \leq A$) or \emptyset (otherwise).)
- ▶ Deduce by Mayer-Vietoris that $\chi_{G,Z}$ is iso if Z has abelian isotropy.
- ▶ Let $F = \{\underline{W} \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$ be the space of complete flags in $\mathbb{C}[G]$, so $Z \times F$ and $Z \times F^2$ have abelian isotropy, and we have projections $p: (Z \times F)_{hG} \rightarrow Z_{hG}$ and $q_0, q_1: (Z \times F^2)_{hG} \rightarrow (Z \times F)_{hG}$
- ▶ We saw before that $E^*((Z \times F)_{hG})$ has a canonical basis $e_1 = 1, e_2, \dots, e_{n!}$ over $E^*(Z_{hG})$. Similarly, the elements $q_0^*(e_i)q_1^*(e_j)$ form a basis for $E^*((Z \times F^2)_{hG})$ over $E^*(Z_{hG})$.
- ▶ It follows that the diagram $E^*(Z_{hG}) \rightarrow E^*((Z \times F)_{hG}) \rightrightarrows E^*((Z \times F^2)_{hG})$ is an equaliser.
- ▶ Deduce the general case from this.

Proof of the generalised character theorem

- ▶ Reduce to the case of a finite group G .
- ▶ Generalise: for a finite G -CW complex Z , we have

$$\chi_{G,Z}: L \otimes_{E^0} E^*(Z_{hG}) \rightarrow L \otimes_{\mathbb{Q}} \left(\prod_{\theta: \Theta^* \rightarrow G} H^*(Z^{\text{image}(\theta)}; \mathbb{Q}) \right)^G$$

- ▶ Prove by calculation that $\theta_{G,Z}$ is iso when $Z = G/A$ with $A \leq G$ abelian. (Here $Z_{hG} = BA$, and $Z^{\text{image}(\theta)}$ is Z (if $\text{image}(\theta) \leq A$) or \emptyset (otherwise).)
- ▶ Deduce by Mayer-Vietoris that $\chi_{G,Z}$ is iso if Z has abelian isotropy.
- ▶ Let $F = \{\underline{W} \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$ be the space of complete flags in $\mathbb{C}[G]$, so $Z \times F$ and $Z \times F^2$ have abelian isotropy, and we have projections $p: (Z \times F)_{hG} \rightarrow Z_{hG}$ and $q_0, q_1: (Z \times F^2)_{hG} \rightarrow (Z \times F)_{hG}$
- ▶ We saw before that $E^*((Z \times F)_{hG})$ has a canonical basis $e_1 = 1, e_2, \dots, e_{n!}$ over $E^*(Z_{hG})$. Similarly, the elements $q_0^*(e_i)q_1^*(e_j)$ form a basis for $E^*((Z \times F^2)_{hG})$ over $E^*(Z_{hG})$.
- ▶ It follows that the diagram $E^*(Z_{hG}) \rightarrow E^*((Z \times F)_{hG}) \rightrightarrows E^*((Z \times F^2)_{hG})$ is an equaliser.
- ▶ Deduce the general case from this.

Proof of the generalised character theorem

- ▶ Reduce to the case of a finite group G .
- ▶ Generalise: for a finite G -CW complex Z , we have

$$\chi_{G,Z}: L \otimes_{E^0} E^*(Z_{hG}) \rightarrow L \otimes_{\mathbb{Q}} \left(\prod_{\theta: \Theta^* \rightarrow G} H^*(Z^{\text{image}(\theta)}; \mathbb{Q}) \right)^G$$

- ▶ Prove by calculation that $\theta_{G,Z}$ is iso when $Z = G/A$ with $A \leq G$ abelian. (Here $Z_{hG} = BA$, and $Z^{\text{image}(\theta)}$ is Z (if $\text{image}(\theta) \leq A$) or \emptyset (otherwise).)
- ▶ Deduce by Mayer-Vietoris that $\chi_{G,Z}$ is iso if Z has abelian isotropy.
- ▶ Let $F = \{ \underline{W} \mid W_0 < \dots < W_n = \mathbb{C}[G] \}$ be the space of complete flags in $\mathbb{C}[G]$, so $Z \times F$ and $Z \times F^2$ have abelian isotropy, and we have projections $p: (Z \times F)_{hG} \rightarrow Z_{hG}$ and $q_0, q_1: (Z \times F^2)_{hG} \rightarrow (Z \times F)_{hG}$
- ▶ We saw before that $E^*((Z \times F)_{hG})$ has a canonical basis $e_1 = 1, e_2, \dots, e_{n!}$ over $E^*(Z_{hG})$. Similarly, the elements $q_0^*(e_i)q_1^*(e_j)$ form a basis for $E^*((Z \times F^2)_{hG})$ over $E^*(Z_{hG})$.
- ▶ It follows that the diagram $E^*(Z_{hG}) \rightarrow E^*((Z \times F)_{hG}) \rightrightarrows E^*((Z \times F^2)_{hG})$ is an equaliser.
- ▶ Deduce the general case from this.

Proof of the generalised character theorem

- ▶ Reduce to the case of a finite group G .
- ▶ Generalise: for a finite G -CW complex Z , we have

$$\chi_{G,Z}: L \otimes_{E^0} E^*(Z_{hG}) \rightarrow L \otimes_{\mathbb{Q}} \left(\prod_{\theta: \Theta^* \rightarrow G} H^*(Z^{\text{image}(\theta)}; \mathbb{Q}) \right)^G$$

- ▶ Prove by calculation that $\theta_{G,Z}$ is iso when $Z = G/A$ with $A \leq G$ abelian. (Here $Z_{hG} = BA$, and $Z^{\text{image}(\theta)}$ is Z (if $\text{image}(\theta) \leq A$) or \emptyset (otherwise).)
- ▶ Deduce by Mayer-Vietoris that $\chi_{G,Z}$ is iso if Z has abelian isotropy.
- ▶ Let $F = \{\underline{W} \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$ be the space of complete flags in $\mathbb{C}[G]$, so $Z \times F$ and $Z \times F^2$ have abelian isotropy, and we have projections $p: (Z \times F)_{hG} \rightarrow Z_{hG}$ and $q_0, q_1: (Z \times F^2)_{hG} \rightarrow (Z \times F)_{hG}$
- ▶ We saw before that $E^*((Z \times F)_{hG})$ has a canonical basis $e_1 = 1, e_2, \dots, e_{n!}$ over $E^*(Z_{hG})$. Similarly, the elements $q_0^*(e_i)q_1^*(e_j)$ form a basis for $E^*((Z \times F^2)_{hG})$ over $E^*(Z_{hG})$.
- ▶ It follows that the diagram $E^*(Z_{hG}) \rightarrow E^*((Z \times F)_{hG}) \rightrightarrows E^*((Z \times F^2)_{hG})$ is an equaliser.
- ▶ Deduce the general case from this.

Proof of the generalised character theorem

- ▶ Reduce to the case of a finite group G .
- ▶ Generalise: for a finite G -CW complex Z , we have

$$\chi_{G,Z}: L \otimes_{E^0} E^*(Z_{hG}) \rightarrow L \otimes_{\mathbb{Q}} \left(\prod_{\theta: \Theta^* \rightarrow G} H^*(Z^{\text{image}(\theta)}; \mathbb{Q}) \right)^G$$

- ▶ Prove by calculation that $\theta_{G,Z}$ is iso when $Z = G/A$ with $A \leq G$ abelian. (Here $Z_{hG} = BA$, and $Z^{\text{image}(\theta)}$ is Z (if $\text{image}(\theta) \leq A$) or \emptyset (otherwise).)
- ▶ Deduce by Mayer-Vietoris that $\chi_{G,Z}$ is iso if Z has abelian isotropy.
- ▶ Let $F = \{\underline{W} \mid W_0 < \dots < W_n = \mathbb{C}[G]\}$ be the space of complete flags in $\mathbb{C}[G]$, so $Z \times F$ and $Z \times F^2$ have abelian isotropy, and we have projections $p: (Z \times F)_{hG} \rightarrow Z_{hG}$ and $q_0, q_1: (Z \times F^2)_{hG} \rightarrow (Z \times F)_{hG}$
- ▶ We saw before that $E^*((Z \times F)_{hG})$ has a canonical basis $e_1 = 1, e_2, \dots, e_{n!}$ over $E^*(Z_{hG})$. Similarly, the elements $q_0^*(e_i)q_1^*(e_j)$ form a basis for $E^*((Z \times F^2)_{hG})$ over $E^*(Z_{hG})$.
- ▶ It follows that the diagram $E^*(Z_{hG}) \rightarrow E^*((Z \times F)_{hG}) \rightrightarrows E^*((Z \times F^2)_{hG})$ is an equaliser.
- ▶ Deduce the general case from this.

Divisors and vector bundles

- ▶ A *divisor of degree d* on $\mathbb{G} = \text{spf}(E^0[[X]])$ is a closed subscheme $D < \mathbb{G}$ such that \mathcal{O}_D is free of rank d as a module over $\mathcal{O}_S = E^0$.
- ▶ Equivalently, $\mathcal{O}_D = E^0[[X]]/f(x)$ for some monic polynomial $f(x) = \sum_{i=0}^d c_i x^{d-i}$ with c_i in the maximal ideal for $i > 0$.
- ▶ $[p^k]_E(x)$ is a unit multiple of a polynomial of degree p^{nk} , so the scheme $\mathbb{G}[p^k] = \ker(p^k \cdot 1: \mathbb{G} \rightarrow \mathbb{G}) = \text{spf}(E^0[[X]]/[p^k]_E(x)) = \text{spf}(E^0(BC_{p^k}))$ is a divisor of degree p^{nk} .
- ▶ More generally, for $T \rightarrow S$, a *divisor of degree d on \mathbb{G} over T* is a closed subscheme $D < T \times_S \mathbb{G}$ such that \mathcal{O}_D is free of rank d over \mathcal{O}_T .
- ▶ Equivalently, $\mathcal{O}_D = \mathcal{O}_T[[X]]/f(x)$ for some monic polynomial $f(x) = \sum_{i=0}^d c_i x^{d-i}$ with $c_i \in \mathcal{O}_T$ topologically nilpotent for $i > 0$.
- ▶ Example: if $V \rightarrow Z$ is a complex bundle of dimension d , then the scheme $D_V = \text{spf}(E^0(P(V)))$ is a divisor of degree d on \mathbb{G} over $\text{spf}(E^0(Z))$ (by earlier calculation of $E^0(P(V))$).
- ▶ There is a sum operation for divisors: if $\mathcal{O}_{D_i} = \mathcal{O}_T[[X]]/f_i(x)$ for $i = 0, 1$ then $\mathcal{O}_{D_0+D_1} = \mathcal{O}_T[[X]]/(f_0(x)f_1(x))$. For this: $D_{V_0 \oplus V_1} = D_{V_0} + D_{V_1}$.
- ▶ An element $a \in \mathbb{G}$ gives a divisor $[a]$ of degree one.
A list a_1, \dots, a_d gives a divisor $\sum_i [a_i]$ of degree d , symmetric in a_1, \dots, a_d .
Using this: $\text{Div}_d^+(\mathbb{G}) = \mathbb{G}^d / \Sigma_d$.

Divisors and vector bundles

- ▶ A *divisor of degree d* on $\mathbb{G} = \text{spf}(E^0[[X]])$ is a closed subscheme $D < \mathbb{G}$ such that \mathcal{O}_D is free of rank d as a module over $\mathcal{O}_S = E^0$.
- ▶ Equivalently, $\mathcal{O}_D = E^0[[X]]/f(x)$ for some monic polynomial $f(x) = \sum_{i=0}^d c_i x^{d-i}$ with c_i in the maximal ideal for $i > 0$.
- ▶ $[p^k]_E(x)$ is a unit multiple of a polynomial of degree p^{nk} , so the scheme $\mathbb{G}[p^k] = \ker(p^k \cdot 1: \mathbb{G} \rightarrow \mathbb{G}) = \text{spf}(E^0[[X]]/[p^k]_E(x)) = \text{spf}(E^0(BC_{p^k}))$ is a divisor of degree p^{nk} .
- ▶ More generally, for $T \rightarrow S$, a *divisor of degree d on \mathbb{G} over T* is a closed subscheme $D < T \times_S \mathbb{G}$ such that \mathcal{O}_D is free of rank d over \mathcal{O}_T .
- ▶ Equivalently, $\mathcal{O}_D = \mathcal{O}_T[[X]]/f(x)$ for some monic polynomial $f(x) = \sum_{i=0}^d c_i x^{d-i}$ with $c_i \in \mathcal{O}_T$ topologically nilpotent for $i > 0$.
- ▶ Example: if $V \rightarrow Z$ is a complex bundle of dimension d , then the scheme $D_V = \text{spf}(E^0(P(V)))$ is a divisor of degree d on \mathbb{G} over $\text{spf}(E^0(Z))$ (by earlier calculation of $E^0(P(V))$).
- ▶ There is a sum operation for divisors: if $\mathcal{O}_{D_i} = \mathcal{O}_T[[X]]/f_i(x)$ for $i = 0, 1$ then $\mathcal{O}_{D_0+D_1} = \mathcal{O}_T[[X]]/(f_0(x)f_1(x))$. For this: $D_{V_0 \oplus V_1} = D_{V_0} + D_{V_1}$.
- ▶ An element $a \in \mathbb{G}$ gives a divisor $[a]$ of degree one.
A list a_1, \dots, a_d gives a divisor $\sum_i [a_i]$ of degree d , symmetric in a_1, \dots, a_d .
Using this: $\text{Div}_d^+(\mathbb{G}) = \mathbb{G}^d / \Sigma_d$.

Divisors and vector bundles

- ▶ A *divisor of degree d* on $\mathbb{G} = \text{spf}(E^0[[X]])$ is a closed subscheme $D < \mathbb{G}$ such that \mathcal{O}_D is free of rank d as a module over $\mathcal{O}_S = E^0$.
- ▶ Equivalently, $\mathcal{O}_D = E^0[[X]]/f(x)$ for some monic polynomial $f(x) = \sum_{i=0}^d c_i x^{d-i}$ with c_i in the maximal ideal for $i > 0$.
- ▶ $[p^k]_E(x)$ is a unit multiple of a polynomial of degree p^{nk} , so the scheme $\mathbb{G}[p^k] = \ker(p^k \cdot 1: \mathbb{G} \rightarrow \mathbb{G}) = \text{spf}(E^0[[X]]/[p^k]_E(x)) = \text{spf}(E^0(BC_{p^k}))$ is a divisor of degree p^{nk} .
- ▶ More generally, for $T \rightarrow S$, a *divisor of degree d on \mathbb{G} over T* is a closed subscheme $D < T \times_S \mathbb{G}$ such that \mathcal{O}_D is free of rank d over \mathcal{O}_T .
- ▶ Equivalently, $\mathcal{O}_D = \mathcal{O}_T[[X]]/f(x)$ for some monic polynomial $f(x) = \sum_{i=0}^d c_i x^{d-i}$ with $c_i \in \mathcal{O}_T$ topologically nilpotent for $i > 0$.
- ▶ Example: if $V \rightarrow Z$ is a complex bundle of dimension d , then the scheme $D_V = \text{spf}(E^0(P(V)))$ is a divisor of degree d on \mathbb{G} over $\text{spf}(E^0(Z))$ (by earlier calculation of $E^0(P(V))$).
- ▶ There is a sum operation for divisors: if $\mathcal{O}_{D_i} = \mathcal{O}_T[[X]]/f_i(x)$ for $i = 0, 1$ then $\mathcal{O}_{D_0+D_1} = \mathcal{O}_T[[X]]/(f_0(x)f_1(x))$. For this: $D_{V_0 \oplus V_1} = D_{V_0} + D_{V_1}$.
- ▶ An element $a \in \mathbb{G}$ gives a divisor $[a]$ of degree one. A list a_1, \dots, a_d gives a divisor $\sum_i [a_i]$ of degree d , symmetric in a_1, \dots, a_d . Using this: $\text{Div}_d^+(\mathbb{G}) = \mathbb{G}^d / \Sigma_d$.

Divisors and vector bundles

- ▶ A *divisor of degree d* on $\mathbb{G} = \text{spf}(E^0[[X]])$ is a closed subscheme $D < \mathbb{G}$ such that \mathcal{O}_D is free of rank d as a module over $\mathcal{O}_S = E^0$.
- ▶ Equivalently, $\mathcal{O}_D = E^0[[X]]/f(x)$ for some monic polynomial $f(x) = \sum_{i=0}^d c_i x^{d-i}$ with c_i in the maximal ideal for $i > 0$.
- ▶ $[p^k]_E(x)$ is a unit multiple of a polynomial of degree p^{nk} , so the scheme $\mathbb{G}[p^k] = \ker(p^k \cdot 1: \mathbb{G} \rightarrow \mathbb{G}) = \text{spf}(E^0[[X]]/[p^k]_E(x)) = \text{spf}(E^0(BC_{p^k}))$ is a divisor of degree p^{nk} .
- ▶ More generally, for $T \rightarrow S$, a *divisor of degree d on \mathbb{G} over T* is a closed subscheme $D < T \times_S \mathbb{G}$ such that \mathcal{O}_D is free of rank d over \mathcal{O}_T .
- ▶ Equivalently, $\mathcal{O}_D = \mathcal{O}_T[[X]]/f(x)$ for some monic polynomial $f(x) = \sum_{i=0}^d c_i x^{d-i}$ with $c_i \in \mathcal{O}_T$ topologically nilpotent for $i > 0$.
- ▶ Example: if $V \rightarrow Z$ is a complex bundle of dimension d , then the scheme $D_V = \text{spf}(E^0(P(V)))$ is a divisor of degree d on \mathbb{G} over $\text{spf}(E^0(Z))$ (by earlier calculation of $E^0(P(V))$).
- ▶ There is a sum operation for divisors: if $\mathcal{O}_{D_i} = \mathcal{O}_T[[X]]/f_i(x)$ for $i = 0, 1$ then $\mathcal{O}_{D_0+D_1} = \mathcal{O}_T[[X]]/(f_0(x)f_1(x))$. For this: $D_{V_0 \oplus V_1} = D_{V_0} + D_{V_1}$.
- ▶ An element $a \in \mathbb{G}$ gives a divisor $[a]$ of degree one. A list a_1, \dots, a_d gives a divisor $\sum_i [a_i]$ of degree d , symmetric in a_1, \dots, a_d . Using this: $\text{Div}_d^+(\mathbb{G}) = \mathbb{G}^d / \Sigma_d$.

Divisors and vector bundles

- ▶ A *divisor of degree d* on $\mathbb{G} = \text{spf}(E^0[[X]])$ is a closed subscheme $D < \mathbb{G}$ such that \mathcal{O}_D is free of rank d as a module over $\mathcal{O}_S = E^0$.
- ▶ Equivalently, $\mathcal{O}_D = E^0[[X]]/f(x)$ for some monic polynomial $f(x) = \sum_{i=0}^d c_i x^{d-i}$ with c_i in the maximal ideal for $i > 0$.
- ▶ $[p^k]_E(x)$ is a unit multiple of a polynomial of degree p^{nk} , so the scheme $\mathbb{G}[p^k] = \ker(p^k \cdot 1: \mathbb{G} \rightarrow \mathbb{G}) = \text{spf}(E^0[[X]]/[p^k]_E(x)) = \text{spf}(E^0(BC_{p^k}))$ is a divisor of degree p^{nk} .
- ▶ More generally, for $T \rightarrow S$, a *divisor of degree d on \mathbb{G} over T* is a closed subscheme $D < T \times_S \mathbb{G}$ such that \mathcal{O}_D is free of rank d over \mathcal{O}_T .
- ▶ Equivalently, $\mathcal{O}_D = \mathcal{O}_T[[X]]/f(x)$ for some monic polynomial $f(x) = \sum_{i=0}^d c_i x^{d-i}$ with $c_i \in \mathcal{O}_T$ topologically nilpotent for $i > 0$.
- ▶ Example: if $V \rightarrow Z$ is a complex bundle of dimension d , then the scheme $D_V = \text{spf}(E^0(P(V)))$ is a divisor of degree d on \mathbb{G} over $\text{spf}(E^0(Z))$ (by earlier calculation of $E^0(P(V))$).
- ▶ There is a sum operation for divisors: if $\mathcal{O}_{D_i} = \mathcal{O}_T[[X]]/f_i(x)$ for $i = 0, 1$ then $\mathcal{O}_{D_0+D_1} = \mathcal{O}_T[[X]]/(f_0(x)f_1(x))$. For this: $D_{V_0 \oplus V_1} = D_{V_0} + D_{V_1}$.
- ▶ An element $a \in \mathbb{G}$ gives a divisor $[a]$ of degree one. A list a_1, \dots, a_d gives a divisor $\sum_i [a_i]$ of degree d , symmetric in a_1, \dots, a_d . Using this: $\text{Div}_d^+(\mathbb{G}) = \mathbb{G}^d / \Sigma_d$.

Divisors and vector bundles

- ▶ A *divisor of degree d* on $\mathbb{G} = \text{spf}(E^0[[X]])$ is a closed subscheme $D < \mathbb{G}$ such that \mathcal{O}_D is free of rank d as a module over $\mathcal{O}_S = E^0$.
- ▶ Equivalently, $\mathcal{O}_D = E^0[[X]]/f(x)$ for some monic polynomial $f(x) = \sum_{i=0}^d c_i x^{d-i}$ with c_i in the maximal ideal for $i > 0$.
- ▶ $[p^k]_E(x)$ is a unit multiple of a polynomial of degree p^{nk} , so the scheme $\mathbb{G}[p^k] = \ker(p^k \cdot 1: \mathbb{G} \rightarrow \mathbb{G}) = \text{spf}(E^0[[X]]/[p^k]_E(x)) = \text{spf}(E^0(BC_{p^k}))$ is a divisor of degree p^{nk} .
- ▶ More generally, for $T \rightarrow S$, a *divisor of degree d on \mathbb{G} over T* is a closed subscheme $D < T \times_S \mathbb{G}$ such that \mathcal{O}_D is free of rank d over \mathcal{O}_T .
- ▶ Equivalently, $\mathcal{O}_D = \mathcal{O}_T[[X]]/f(x)$ for some monic polynomial $f(x) = \sum_{i=0}^d c_i x^{d-i}$ with $c_i \in \mathcal{O}_T$ topologically nilpotent for $i > 0$.
- ▶ Example: if $V \rightarrow Z$ is a complex bundle of dimension d , then the scheme $D_V = \text{spf}(E^0(P(V)))$ is a divisor of degree d on \mathbb{G} over $\text{spf}(E^0(Z))$ (by earlier calculation of $E^0(P(V))$).
- ▶ There is a sum operation for divisors: if $\mathcal{O}_{D_i} = \mathcal{O}_T[[X]]/f_i(x)$ for $i = 0, 1$ then $\mathcal{O}_{D_0+D_1} = \mathcal{O}_T[[X]]/(f_0(x)f_1(x))$. For this: $D_{V_0 \oplus V_1} = D_{V_0} + D_{V_1}$.
- ▶ An element $a \in \mathbb{G}$ gives a divisor $[a]$ of degree one. A list a_1, \dots, a_d gives a divisor $\sum_i [a_i]$ of degree d , symmetric in a_1, \dots, a_d . Using this: $\text{Div}_d^+(\mathbb{G}) = \mathbb{G}^d / \Sigma_d$.

Divisors and vector bundles

- ▶ A *divisor of degree d* on $\mathbb{G} = \text{spf}(E^0[[X]])$ is a closed subscheme $D < \mathbb{G}$ such that \mathcal{O}_D is free of rank d as a module over $\mathcal{O}_S = E^0$.
- ▶ Equivalently, $\mathcal{O}_D = E^0[[X]]/f(x)$ for some monic polynomial $f(x) = \sum_{i=0}^d c_i x^{d-i}$ with c_i in the maximal ideal for $i > 0$.
- ▶ $[p^k]_E(x)$ is a unit multiple of a polynomial of degree p^{nk} , so the scheme $\mathbb{G}[p^k] = \ker(p^k \cdot 1: \mathbb{G} \rightarrow \mathbb{G}) = \text{spf}(E^0[[X]]/[p^k]_E(x)) = \text{spf}(E^0(BC_{p^k}))$ is a divisor of degree p^{nk} .
- ▶ More generally, for $T \rightarrow S$, a *divisor of degree d on \mathbb{G} over T* is a closed subscheme $D < T \times_S \mathbb{G}$ such that \mathcal{O}_D is free of rank d over \mathcal{O}_T .
- ▶ Equivalently, $\mathcal{O}_D = \mathcal{O}_T[[X]]/f(x)$ for some monic polynomial $f(x) = \sum_{i=0}^d c_i x^{d-i}$ with $c_i \in \mathcal{O}_T$ topologically nilpotent for $i > 0$.
- ▶ Example: if $V \rightarrow Z$ is a complex bundle of dimension d , then the scheme $D_V = \text{spf}(E^0(P(V)))$ is a divisor of degree d on \mathbb{G} over $\text{spf}(E^0(Z))$ (by earlier calculation of $E^0(P(V))$).
- ▶ There is a sum operation for divisors: if $\mathcal{O}_{D_i} = \mathcal{O}_T[[X]]/f_i(x)$ for $i = 0, 1$ then $\mathcal{O}_{D_0+D_1} = \mathcal{O}_T[[X]]/(f_0(x)f_1(x))$. For this: $D_{V_0 \oplus V_1} = D_{V_0} + D_{V_1}$.
- ▶ An element $a \in \mathbb{G}$ gives a divisor $[a]$ of degree one.
A list a_1, \dots, a_d gives a divisor $\sum_i [a_i]$ of degree d , symmetric in a_1, \dots, a_d .
Using this: $\text{Div}_d^+(\mathbb{G}) = \mathbb{G}^d / \Sigma_d$.

Divisors and vector bundles

- ▶ A *divisor of degree d* on $\mathbb{G} = \text{spf}(E^0[[X]])$ is a closed subscheme $D < \mathbb{G}$ such that \mathcal{O}_D is free of rank d as a module over $\mathcal{O}_S = E^0$.
- ▶ Equivalently, $\mathcal{O}_D = E^0[[X]]/f(x)$ for some monic polynomial $f(x) = \sum_{i=0}^d c_i x^{d-i}$ with c_i in the maximal ideal for $i > 0$.
- ▶ $[p^k]_E(x)$ is a unit multiple of a polynomial of degree p^{nk} , so the scheme $\mathbb{G}[p^k] = \ker(p^k \cdot 1: \mathbb{G} \rightarrow \mathbb{G}) = \text{spf}(E^0[[X]]/[p^k]_E(x)) = \text{spf}(E^0(BC_{p^k}))$ is a divisor of degree p^{nk} .
- ▶ More generally, for $T \rightarrow S$, a *divisor of degree d on \mathbb{G} over T* is a closed subscheme $D < T \times_S \mathbb{G}$ such that \mathcal{O}_D is free of rank d over \mathcal{O}_T .
- ▶ Equivalently, $\mathcal{O}_D = \mathcal{O}_T[[X]]/f(x)$ for some monic polynomial $f(x) = \sum_{i=0}^d c_i x^{d-i}$ with $c_i \in \mathcal{O}_T$ topologically nilpotent for $i > 0$.
- ▶ Example: if $V \rightarrow Z$ is a complex bundle of dimension d , then the scheme $D_V = \text{spf}(E^0(P(V)))$ is a divisor of degree d on \mathbb{G} over $\text{spf}(E^0(Z))$ (by earlier calculation of $E^0(P(V))$).
- ▶ There is a sum operation for divisors: if $\mathcal{O}_{D_i} = \mathcal{O}_T[[X]]/f_i(x)$ for $i = 0, 1$ then $\mathcal{O}_{D_0+D_1} = \mathcal{O}_T[[X]]/(f_0(x)f_1(x))$. For this: $D_{V_0 \oplus V_1} = D_{V_0} + D_{V_1}$.
- ▶ An element $a \in \mathbb{G}$ gives a divisor $[a]$ of degree one.
A list a_1, \dots, a_d gives a divisor $\sum_i [a_i]$ of degree d , symmetric in a_1, \dots, a_d .
Using this: $\text{Div}_d^+(\mathbb{G}) = \mathbb{G}^d / \Sigma_d$.

Divisors and vector bundles

- ▶ A *divisor of degree d* on $\mathbb{G} = \text{spf}(E^0[[X]])$ is a closed subscheme $D < \mathbb{G}$ such that \mathcal{O}_D is free of rank d as a module over $\mathcal{O}_S = E^0$.
- ▶ Equivalently, $\mathcal{O}_D = E^0[[X]]/f(x)$ for some monic polynomial $f(x) = \sum_{i=0}^d c_i x^{d-i}$ with c_i in the maximal ideal for $i > 0$.
- ▶ $[p^k]_E(x)$ is a unit multiple of a polynomial of degree p^{nk} , so the scheme $\mathbb{G}[p^k] = \ker(p^k \cdot 1: \mathbb{G} \rightarrow \mathbb{G}) = \text{spf}(E^0[[X]]/[p^k]_E(x)) = \text{spf}(E^0(BC_{p^k}))$ is a divisor of degree p^{nk} .
- ▶ More generally, for $T \rightarrow S$, a *divisor of degree d on \mathbb{G} over T* is a closed subscheme $D < T \times_S \mathbb{G}$ such that \mathcal{O}_D is free of rank d over \mathcal{O}_T .
- ▶ Equivalently, $\mathcal{O}_D = \mathcal{O}_T[[X]]/f(x)$ for some monic polynomial $f(x) = \sum_{i=0}^d c_i x^{d-i}$ with $c_i \in \mathcal{O}_T$ topologically nilpotent for $i > 0$.
- ▶ Example: if $V \rightarrow Z$ is a complex bundle of dimension d , then the scheme $D_V = \text{spf}(E^0(P(V)))$ is a divisor of degree d on \mathbb{G} over $\text{spf}(E^0(Z))$ (by earlier calculation of $E^0(P(V))$).
- ▶ There is a sum operation for divisors: if $\mathcal{O}_{D_i} = \mathcal{O}_T[[X]]/f_i(x)$ for $i = 0, 1$ then $\mathcal{O}_{D_0+D_1} = \mathcal{O}_T[[X]]/(f_0(x)f_1(x))$. For this: $D_{V_0 \oplus V_1} = D_{V_0} + D_{V_1}$.
- ▶ An element $a \in \mathbb{G}$ gives a divisor $[a]$ of degree one.
A list a_1, \dots, a_d gives a divisor $\sum_i [a_i]$ of degree d , symmetric in a_1, \dots, a_d .
Using this: $\text{Div}_d^+(\mathbb{G}) = \mathbb{G}^d / \Sigma_d$.

Divisors and vector bundles

- ▶ A *divisor of degree d* on $\mathbb{G} = \text{spf}(E^0[[X]])$ is a closed subscheme $D < \mathbb{G}$ such that \mathcal{O}_D is free of rank d as a module over $\mathcal{O}_S = E^0$.
- ▶ Equivalently, $\mathcal{O}_D = E^0[[X]]/f(x)$ for some monic polynomial $f(x) = \sum_{i=0}^d c_i x^{d-i}$ with c_i in the maximal ideal for $i > 0$.
- ▶ $[p^k]_E(x)$ is a unit multiple of a polynomial of degree p^{nk} , so the scheme $\mathbb{G}[p^k] = \ker(p^k \cdot 1: \mathbb{G} \rightarrow \mathbb{G}) = \text{spf}(E^0[[X]]/[p^k]_E(x)) = \text{spf}(E^0(BC_{p^k}))$ is a divisor of degree p^{nk} .
- ▶ More generally, for $T \rightarrow S$, a *divisor of degree d on \mathbb{G} over T* is a closed subscheme $D < T \times_S \mathbb{G}$ such that \mathcal{O}_D is free of rank d over \mathcal{O}_T .
- ▶ Equivalently, $\mathcal{O}_D = \mathcal{O}_T[[X]]/f(x)$ for some monic polynomial $f(x) = \sum_{i=0}^d c_i x^{d-i}$ with $c_i \in \mathcal{O}_T$ topologically nilpotent for $i > 0$.
- ▶ Example: if $V \rightarrow Z$ is a complex bundle of dimension d , then the scheme $D_V = \text{spf}(E^0(P(V)))$ is a divisor of degree d on \mathbb{G} over $\text{spf}(E^0(Z))$ (by earlier calculation of $E^0(P(V))$).
- ▶ There is a sum operation for divisors: if $\mathcal{O}_{D_i} = \mathcal{O}_T[[X]]/f_i(x)$ for $i = 0, 1$ then $\mathcal{O}_{D_0+D_1} = \mathcal{O}_T[[X]]/(f_0(x)f_1(x))$. For this: $D_{V_0 \oplus V_1} = D_{V_0} + D_{V_1}$.
- ▶ An element $a \in \mathbb{G}$ gives a divisor $[a]$ of degree one. A list a_1, \dots, a_d gives a divisor $\sum_i [a_i]$ of degree d , symmetric in a_1, \dots, a_d . Using this: $\text{Div}_d^+(\mathbb{G}) = \mathbb{G}^d / \Sigma_d$.

- ▶ A *divisor of degree d* on $\mathbb{G} = \text{spf}(E^0[[X]])$ is a closed subscheme $D < \mathbb{G}$ such that \mathcal{O}_D is free of rank d as a module over $\mathcal{O}_S = E^0$.
- ▶ Equivalently, $\mathcal{O}_D = E^0[[X]]/f(x)$ for some monic polynomial $f(x) = \sum_{i=0}^d c_i x^{d-i}$ with c_i in the maximal ideal for $i > 0$.
- ▶ $[p^k]_E(x)$ is a unit multiple of a polynomial of degree p^{nk} , so the scheme $\mathbb{G}[p^k] = \ker(p^k \cdot 1: \mathbb{G} \rightarrow \mathbb{G}) = \text{spf}(E^0[[X]]/[p^k]_E(x)) = \text{spf}(E^0(BC_{p^k}))$ is a divisor of degree p^{nk} .
- ▶ More generally, for $T \rightarrow S$, a *divisor of degree d on \mathbb{G} over T* is a closed subscheme $D < T \times_S \mathbb{G}$ such that \mathcal{O}_D is free of rank d over \mathcal{O}_T .
- ▶ Equivalently, $\mathcal{O}_D = \mathcal{O}_T[[X]]/f(x)$ for some monic polynomial $f(x) = \sum_{i=0}^d c_i x^{d-i}$ with $c_i \in \mathcal{O}_T$ topologically nilpotent for $i > 0$.
- ▶ Example: if $V \rightarrow Z$ is a complex bundle of dimension d , then the scheme $D_V = \text{spf}(E^0(P(V)))$ is a divisor of degree d on \mathbb{G} over $\text{spf}(E^0(Z))$ (by earlier calculation of $E^0(P(V))$).
- ▶ There is a sum operation for divisors: if $\mathcal{O}_{D_i} = \mathcal{O}_T[[X]]/f_i(x)$ for $i = 0, 1$ then $\mathcal{O}_{D_0+D_1} = \mathcal{O}_T[[X]]/(f_0(x)f_1(x))$. For this: $D_{V_0 \oplus V_1} = D_{V_0} + D_{V_1}$.
- ▶ An element $a \in \mathbb{G}$ gives a divisor $[a]$ of degree one. A list a_1, \dots, a_d gives a divisor $\sum_i [a_i]$ of degree d , symmetric in a_1, \dots, a_d . Using this: $\text{Div}_d^+(\mathbb{G}) = \mathbb{G}^d / \Sigma_d$.

- ▶ We have identified $\mathrm{spf}(E^0(BA))$ with $\mathrm{Hom}(A^*, \mathbb{G})$.
- ▶ Is there a subscheme of monomorphisms from A^* to \mathbb{G} ?
- ▶ Monomorphisms of schemes are closely related to epimorphisms of commutative rings, which have poor behaviour. Note that $\mathbb{Z} \rightarrow \mathbb{Q}$ and $\mathbb{Z} \rightarrow \mathbb{Z}/n$ are epimorphisms, but behave quite differently.
- ▶ We will define $\mathrm{Level}(A^*, \mathbb{G}) \subseteq \mathrm{Hom}(A^*, \mathbb{G})$ which is motivated by the above; but do not take the analogy too seriously.
- ▶ Write $A = \prod_{i=0}^{s-1} C_{p^{m_i+1}}$, put $D'_A = E^0[[x_i \mid i < s]]$ and $y_i = [p^{m_i}](x_i) \in R'$.
- ▶ Let U be the set of all terms $\sum_{i < s}^F [k_i]_E(y_i)$ with $0 \leq k_i < p$ for all i .
- ▶ Put $g(t) = \prod_{u \in U} (t - u)$ and $[p]_E(t) = q(t)g(t) + r(t)$ with $\deg(r(t)) < s$. Let I be the ideal generated by the coefficients of $r(t)$, and $D_A = D'_A/I$, and $\mathrm{Level}(A^*, \mathbb{G}) = \mathrm{spf}(D_A)$.
- ▶ Schematically:

$$\mathrm{Level}(A^*, \mathbb{G}) = \{ \phi \in \mathrm{Hom}(A^*, \mathbb{G}) \mid \sum_{\alpha \in A^*[p]} [\phi(\alpha)] \leq \mathbb{G}[p] \}.$$
- ▶ From HKR: $\mathbb{Q} \otimes E^0(B\mathbb{G}) = u_0^{-1} E^0(B\mathbb{G}) \simeq \left(\prod_{A \leq \mathbb{G}} \mathbb{Q} \otimes D_A \right)^{\mathbb{G}}$.
- ▶ There is a similar map $u_k^{-1} E^0(B\mathbb{G})/I_k \rightarrow \left(\prod_{A \leq \mathbb{G}} u_k^{-1} D_{k,A} \right)^{\mathbb{G}}$ for $k > 0$, which is an F -isomorphism (Greenlees-Strickland; see also Stapleton).

- ▶ We have identified $\mathrm{spf}(E^0(BA))$ with $\mathrm{Hom}(A^*, \mathbb{G})$.
- ▶ Is there a subscheme of monomorphisms from A^* to \mathbb{G} ?
- ▶ Monomorphisms of schemes are closely related to epimorphisms of commutative rings, which have poor behaviour. Note that $\mathbb{Z} \rightarrow \mathbb{Q}$ and $\mathbb{Z} \rightarrow \mathbb{Z}/n$ are epimorphisms, but behave quite differently.
- ▶ We will define $\mathrm{Level}(A^*, \mathbb{G}) \subseteq \mathrm{Hom}(A^*, \mathbb{G})$ which is motivated by the above; but do not take the analogy too seriously.
- ▶ Write $A = \prod_{i=0}^{s-1} C_{p^{m_i+1}}$, put $D'_A = E^0[[x_i \mid i < s]]$ and $y_i = [p^{m_i}](x_i) \in R'$.
- ▶ Let U be the set of all terms $\sum_{i < s}^F [k_i]_E(y_i)$ with $0 \leq k_i < p$ for all i .
- ▶ Put $g(t) = \prod_{u \in U} (t - u)$ and $[p]_E(t) = q(t)g(t) + r(t)$ with $\deg(r(t)) < s$. Let I be the ideal generated by the coefficients of $r(t)$, and $D_A = D'_A/I$, and $\mathrm{Level}(A^*, \mathbb{G}) = \mathrm{spf}(D_A)$.
- ▶ Schematically:

$$\mathrm{Level}(A^*, \mathbb{G}) = \{ \phi \in \mathrm{Hom}(A^*, \mathbb{G}) \mid \sum_{\alpha \in A^*[p]} [\phi(\alpha)] \leq \mathbb{G}[p] \}.$$
- ▶ From HKR: $\mathbb{Q} \otimes E^0(BG) = u_0^{-1} E^0(BG) \simeq \left(\prod_{A \leq G} \mathbb{Q} \otimes D_A \right)^G$.
- ▶ There is a similar map $u_k^{-1} E^0(BG)/I_k \rightarrow \left(\prod_{A \leq G} u_k^{-1} D_{k,A} \right)^G$ for $k > 0$, which is an F -isomorphism (Greenlees-Strickland; see also Stapleton).

- ▶ We have identified $\mathrm{spf}(E^0(BA))$ with $\mathrm{Hom}(A^*, \mathbb{G})$.
- ▶ Is there a subscheme of monomorphisms from A^* to \mathbb{G} ?
- ▶ Monomorphisms of schemes are closely related to epimorphisms of commutative rings, which have poor behaviour. Note that $\mathbb{Z} \rightarrow \mathbb{Q}$ and $\mathbb{Z} \rightarrow \mathbb{Z}/n$ are epimorphisms, but behave quite differently.
- ▶ We will define $\mathrm{Level}(A^*, \mathbb{G}) \subseteq \mathrm{Hom}(A^*, \mathbb{G})$ which is motivated by the above; but do not take the analogy too seriously.
- ▶ Write $A = \prod_{i=0}^{s-1} C_{p^{m_i+1}}$, put $D'_A = E^0[[x_i \mid i < s]]$ and $y_i = [p^{m_i}](x_i) \in R'$.
- ▶ Let U be the set of all terms $\sum_{i < s}^F [k_i]_E(y_i)$ with $0 \leq k_i < p$ for all i .
- ▶ Put $g(t) = \prod_{u \in U} (t - u)$ and $[p]_E(t) = q(t)g(t) + r(t)$ with $\deg(r(t)) < s$. Let I be the ideal generated by the coefficients of $r(t)$, and $D_A = D'_A/I$, and $\mathrm{Level}(A^*, \mathbb{G}) = \mathrm{spf}(D_A)$.
- ▶ Schematically:

$$\mathrm{Level}(A^*, \mathbb{G}) = \{ \phi \in \mathrm{Hom}(A^*, \mathbb{G}) \mid \sum_{\alpha \in A^*[p]} [\phi(\alpha)] \leq \mathbb{G}[p] \}.$$
- ▶ From HKR: $\mathbb{Q} \otimes E^0(BG) = u_0^{-1} E^0(BG) \simeq \left(\prod_{A \leq G} \mathbb{Q} \otimes D_A \right)^G$.
- ▶ There is a similar map $u_k^{-1} E^0(BG)/I_k \rightarrow \left(\prod_{A \leq G} u_k^{-1} D_{k,A} \right)^G$ for $k > 0$, which is an F -isomorphism (Greenlees-Strickland; see also Stapleton).

- ▶ We have identified $\mathrm{spf}(E^0(BA))$ with $\mathrm{Hom}(A^*, \mathbb{G})$.
- ▶ Is there a subscheme of monomorphisms from A^* to \mathbb{G} ?
- ▶ Monomorphisms of schemes are closely related to epimorphisms of commutative rings, which have poor behaviour. Note that $\mathbb{Z} \rightarrow \mathbb{Q}$ and $\mathbb{Z} \rightarrow \mathbb{Z}/n$ are epimorphisms, but behave quite differently.
- ▶ We will define $\mathrm{Level}(A^*, \mathbb{G}) \subseteq \mathrm{Hom}(A^*, \mathbb{G})$ which is motivated by the above; but do not take the analogy too seriously.
- ▶ Write $A = \prod_{i=0}^{s-1} C_{p^{m_i+1}}$, put $D'_A = E^0[[x_i \mid i < s]]$ and $y_i = [p^{m_i}](x_i) \in R'$.
- ▶ Let U be the set of all terms $\sum_{i < s}^F [k_i]_E(y_i)$ with $0 \leq k_i < p$ for all i .
- ▶ Put $g(t) = \prod_{u \in U} (t - u)$ and $[p]_E(t) = q(t)g(t) + r(t)$ with $\deg(r(t)) < s$. Let I be the ideal generated by the coefficients of $r(t)$, and $D_A = D'_A/I$, and $\mathrm{Level}(A^*, \mathbb{G}) = \mathrm{spf}(D_A)$.
- ▶ Schematically:

$$\mathrm{Level}(A^*, \mathbb{G}) = \{\phi \in \mathrm{Hom}(A^*, \mathbb{G}) \mid \sum_{\alpha \in A^*[p]} [\phi(\alpha)] \leq \mathbb{G}[p]\}.$$
- ▶ From HKR: $\mathbb{Q} \otimes E^0(BG) = u_0^{-1} E^0(BG) \simeq \left(\prod_{A \leq G} \mathbb{Q} \otimes D_A \right)^G$.
- ▶ There is a similar map $u_k^{-1} E^0(BG)/I_k \rightarrow \left(\prod_{A \leq G} u_k^{-1} D_{k,A} \right)^G$ for $k > 0$, which is an F -isomorphism (Greenlees-Strickland; see also Stapleton).

- ▶ We have identified $\mathrm{spf}(E^0(BA))$ with $\mathrm{Hom}(A^*, \mathbb{G})$.
- ▶ Is there a subscheme of monomorphisms from A^* to \mathbb{G} ?
- ▶ Monomorphisms of schemes are closely related to epimorphisms of commutative rings, which have poor behaviour. Note that $\mathbb{Z} \rightarrow \mathbb{Q}$ and $\mathbb{Z} \rightarrow \mathbb{Z}/n$ are epimorphisms, but behave quite differently.
- ▶ We will define $\mathrm{Level}(A^*, \mathbb{G}) \subseteq \mathrm{Hom}(A^*, \mathbb{G})$ which is motivated by the above; but do not take the analogy too seriously.
- ▶ Write $A = \prod_{i=0}^{s-1} C_{p^{m_i+1}}$, put $D'_A = E^0[[x_i \mid i < s]]$ and $y_i = [p^{m_i}](x_i) \in R'$.
- ▶ Let U be the set of all terms $\sum_{i < s}^F [k_i]_E(y_i)$ with $0 \leq k_i < p$ for all i .
- ▶ Put $g(t) = \prod_{u \in U} (t - u)$ and $[p]_E(t) = q(t)g(t) + r(t)$ with $\deg(r(t)) < s$. Let I be the ideal generated by the coefficients of $r(t)$, and $D_A = D'_A/I$, and $\mathrm{Level}(A^*, \mathbb{G}) = \mathrm{spf}(D_A)$.
- ▶ Schematically:

$$\mathrm{Level}(A^*, \mathbb{G}) = \{ \phi \in \mathrm{Hom}(A^*, \mathbb{G}) \mid \sum_{\alpha \in A^*[p]} [\phi(\alpha)] \leq \mathbb{G}[p] \}.$$
- ▶ From HKR: $\mathbb{Q} \otimes E^0(BG) = u_0^{-1} E^0(BG) \simeq \left(\prod_{A \leq G} \mathbb{Q} \otimes D_A \right)^G$.
- ▶ There is a similar map $u_k^{-1} E^0(BG)/I_k \rightarrow \left(\prod_{A \leq G} u_k^{-1} D_{k,A} \right)^G$ for $k > 0$, which is an F -isomorphism (Greenlees-Strickland; see also Stapleton).

- ▶ We have identified $\mathrm{spf}(E^0(BA))$ with $\mathrm{Hom}(A^*, \mathbb{G})$.
- ▶ Is there a subscheme of monomorphisms from A^* to \mathbb{G} ?
- ▶ Monomorphisms of schemes are closely related to epimorphisms of commutative rings, which have poor behaviour. Note that $\mathbb{Z} \rightarrow \mathbb{Q}$ and $\mathbb{Z} \rightarrow \mathbb{Z}/n$ are epimorphisms, but behave quite differently.
- ▶ We will define $\mathrm{Level}(A^*, \mathbb{G}) \subseteq \mathrm{Hom}(A^*, \mathbb{G})$ which is motivated by the above; but do not take the analogy too seriously.
- ▶ Write $A = \prod_{i=0}^{s-1} C_{p^{m_i+1}}$, put $D'_A = E^0[[x_i \mid i < s]]$ and $y_i = [p^{m_i}](x_i) \in R'$.
- ▶ Let U be the set of all terms $\sum_{i < s}^F [k_i]_E(y_i)$ with $0 \leq k_i < p$ for all i .
- ▶ Put $g(t) = \prod_{u \in U} (t - u)$ and $[p]_E(t) = q(t)g(t) + r(t)$ with $\deg(r(t)) < s$. Let I be the ideal generated by the coefficients of $r(t)$, and $D_A = D'_A/I$, and $\mathrm{Level}(A^*, \mathbb{G}) = \mathrm{spf}(D_A)$.
- ▶ Schematically:

$$\mathrm{Level}(A^*, \mathbb{G}) = \{ \phi \in \mathrm{Hom}(A^*, \mathbb{G}) \mid \sum_{\alpha \in A^*[p]} [\phi(\alpha)] \leq \mathbb{G}[p] \}.$$
- ▶ From HKR: $\mathbb{Q} \otimes E^0(BG) = u_0^{-1} E^0(BG) \simeq \left(\prod_{A \leq G} \mathbb{Q} \otimes D_A \right)^G$.
- ▶ There is a similar map $u_k^{-1} E^0(BG)/I_k \rightarrow \left(\prod_{A \leq G} u_k^{-1} D_{k,A} \right)^G$ for $k > 0$, which is an F -isomorphism (Greenlees-Strickland; see also Stapleton).

- ▶ We have identified $\mathrm{spf}(E^0(BA))$ with $\mathrm{Hom}(A^*, \mathbb{G})$.
- ▶ Is there a subscheme of monomorphisms from A^* to \mathbb{G} ?
- ▶ Monomorphisms of schemes are closely related to epimorphisms of commutative rings, which have poor behaviour. Note that $\mathbb{Z} \rightarrow \mathbb{Q}$ and $\mathbb{Z} \rightarrow \mathbb{Z}/n$ are epimorphisms, but behave quite differently.
- ▶ We will define $\mathrm{Level}(A^*, \mathbb{G}) \subseteq \mathrm{Hom}(A^*, \mathbb{G})$ which is motivated by the above; but do not take the analogy too seriously.
- ▶ Write $A = \prod_{i=0}^{s-1} C_{p^{m_i+1}}$, put $D'_A = E^0[[x_i \mid i < s]]$ and $y_i = [p^{m_i}](x_i) \in R'$.
- ▶ Let U be the set of all terms $\sum_{i < s}^F [k_i]_E(y_i)$ with $0 \leq k_i < p$ for all i .
- ▶ Put $g(t) = \prod_{u \in U} (t - u)$ and $[p]_E(t) = q(t)g(t) + r(t)$ with $\deg(r(t)) < s$. Let I be the ideal generated by the coefficients of $r(t)$, and $D_A = D'_A/I$, and $\mathrm{Level}(A^*, \mathbb{G}) = \mathrm{spf}(D_A)$.
- ▶ Schematically:

$$\mathrm{Level}(A^*, \mathbb{G}) = \{ \phi \in \mathrm{Hom}(A^*, \mathbb{G}) \mid \sum_{\alpha \in A^*[p]} [\phi(\alpha)] \leq \mathbb{G}[p] \}.$$
- ▶ From HKR: $\mathbb{Q} \otimes E^0(BG) = u_0^{-1} E^0(BG) \simeq \left(\prod_{A \leq G} \mathbb{Q} \otimes D_A \right)^G$.
- ▶ There is a similar map $u_k^{-1} E^0(BG)/I_k \rightarrow \left(\prod_{A \leq G} u_k^{-1} D_{k,A} \right)^G$ for $k > 0$, which is an F -isomorphism (Greenlees-Strickland; see also Stapleton).

- ▶ We have identified $\mathrm{spf}(E^0(BA))$ with $\mathrm{Hom}(A^*, \mathbb{G})$.
- ▶ Is there a subscheme of monomorphisms from A^* to \mathbb{G} ?
- ▶ Monomorphisms of schemes are closely related to epimorphisms of commutative rings, which have poor behaviour. Note that $\mathbb{Z} \rightarrow \mathbb{Q}$ and $\mathbb{Z} \rightarrow \mathbb{Z}/n$ are epimorphisms, but behave quite differently.
- ▶ We will define $\mathrm{Level}(A^*, \mathbb{G}) \subseteq \mathrm{Hom}(A^*, \mathbb{G})$ which is motivated by the above; but do not take the analogy too seriously.
- ▶ Write $A = \prod_{i=0}^{s-1} C_{p^{m_i+1}}$, put $D'_A = E^0[[x_i \mid i < s]]$ and $y_i = [p^{m_i}](x_i) \in R'$.
- ▶ Let U be the set of all terms $\sum_{i < s}^F [k_i]_E(y_i)$ with $0 \leq k_i < p$ for all i .
- ▶ Put $g(t) = \prod_{u \in U} (t - u)$ and $[p]_E(t) = q(t)g(t) + r(t)$ with $\deg(r(t)) < s$. Let I be the ideal generated by the coefficients of $r(t)$, and $D_A = D'_A/I$, and $\mathrm{Level}(A^*, \mathbb{G}) = \mathrm{spf}(D_A)$.
- ▶ Schematically:

$$\mathrm{Level}(A^*, \mathbb{G}) = \{ \phi \in \mathrm{Hom}(A^*, \mathbb{G}) \mid \sum_{\alpha \in A^*[p]} [\phi(\alpha)] \leq \mathbb{G}[p] \}.$$
- ▶ From HKR: $\mathbb{Q} \otimes E^0(B\mathbb{G}) = u_0^{-1} E^0(B\mathbb{G}) \simeq \left(\prod_{A \leq \mathbb{G}} \mathbb{Q} \otimes D_A \right)^{\mathbb{G}}$.
- ▶ There is a similar map $u_k^{-1} E^0(B\mathbb{G})/I_k \rightarrow \left(\prod_{A \leq \mathbb{G}} u_k^{-1} D_{k,A} \right)^{\mathbb{G}}$ for $k > 0$, which is an F -isomorphism (Greenlees-Strickland; see also Stapleton).

- ▶ We have identified $\mathrm{spf}(E^0(BA))$ with $\mathrm{Hom}(A^*, \mathbb{G})$.
- ▶ Is there a subscheme of monomorphisms from A^* to \mathbb{G} ?
- ▶ Monomorphisms of schemes are closely related to epimorphisms of commutative rings, which have poor behaviour. Note that $\mathbb{Z} \rightarrow \mathbb{Q}$ and $\mathbb{Z} \rightarrow \mathbb{Z}/n$ are epimorphisms, but behave quite differently.
- ▶ We will define $\mathrm{Level}(A^*, \mathbb{G}) \subseteq \mathrm{Hom}(A^*, \mathbb{G})$ which is motivated by the above; but do not take the analogy too seriously.
- ▶ Write $A = \prod_{i=0}^{s-1} C_{p^{m_i+1}}$, put $D'_A = E^0[[x_i \mid i < s]]$ and $y_i = [p^{m_i}](x_i) \in R'$.
- ▶ Let U be the set of all terms $\sum_{i < s}^F [k_i]_E(y_i)$ with $0 \leq k_i < p$ for all i .
- ▶ Put $g(t) = \prod_{u \in U} (t - u)$ and $[p]_E(t) = q(t)g(t) + r(t)$ with $\deg(r(t)) < s$. Let I be the ideal generated by the coefficients of $r(t)$, and $D_A = D'_A/I$, and $\mathrm{Level}(A^*, \mathbb{G}) = \mathrm{spf}(D_A)$.
- ▶ Schematically:

$$\mathrm{Level}(A^*, \mathbb{G}) = \{\phi \in \mathrm{Hom}(A^*, \mathbb{G}) \mid \sum_{\alpha \in A^*[\rho]} [\phi(\alpha)] \leq \mathbb{G}[\rho]\}.$$
- ▶ From HKR: $\mathbb{Q} \otimes E^0(B\mathbb{G}) = u_0^{-1} E^0(B\mathbb{G}) \simeq \left(\prod_{A \leq \mathbb{G}} \mathbb{Q} \otimes D_A \right)^{\mathbb{G}}$.
- ▶ There is a similar map $u_k^{-1} E^0(B\mathbb{G})/I_k \rightarrow \left(\prod_{A \leq \mathbb{G}} u_k^{-1} D_{k,A} \right)^{\mathbb{G}}$ for $k > 0$, which is an F -isomorphism (Greenlees-Strickland; see also Stapleton).

- ▶ We have identified $\mathrm{spf}(E^0(BA))$ with $\mathrm{Hom}(A^*, \mathbb{G})$.
- ▶ Is there a subscheme of monomorphisms from A^* to \mathbb{G} ?
- ▶ Monomorphisms of schemes are closely related to epimorphisms of commutative rings, which have poor behaviour. Note that $\mathbb{Z} \rightarrow \mathbb{Q}$ and $\mathbb{Z} \rightarrow \mathbb{Z}/n$ are epimorphisms, but behave quite differently.
- ▶ We will define $\mathrm{Level}(A^*, \mathbb{G}) \subseteq \mathrm{Hom}(A^*, \mathbb{G})$ which is motivated by the above; but do not take the analogy too seriously.
- ▶ Write $A = \prod_{i=0}^{s-1} C_{p^{m_i+1}}$, put $D'_A = E^0[[x_i \mid i < s]]$ and $y_i = [p^{m_i}](x_i) \in R'$.
- ▶ Let U be the set of all terms $\sum_{i < s}^F [k_i]_E(y_i)$ with $0 \leq k_i < p$ for all i .
- ▶ Put $g(t) = \prod_{u \in U} (t - u)$ and $[p]_E(t) = q(t)g(t) + r(t)$ with $\deg(r(t)) < s$. Let I be the ideal generated by the coefficients of $r(t)$, and $D_A = D'_A/I$, and $\mathrm{Level}(A^*, \mathbb{G}) = \mathrm{spf}(D_A)$.
- ▶ Schematically:

$$\mathrm{Level}(A^*, \mathbb{G}) = \{\phi \in \mathrm{Hom}(A^*, \mathbb{G}) \mid \sum_{\alpha \in A^*[\rho]} [\phi(\alpha)] \leq \mathbb{G}[\rho]\}.$$
- ▶ From HKR: $\mathbb{Q} \otimes E^0(BG) = u_0^{-1} E^0(BG) \simeq \left(\prod_{A \leq G} \mathbb{Q} \otimes D_A \right)^G$.
- ▶ There is a similar map $u_k^{-1} E^0(BG)/I_k \rightarrow \left(\prod_{A \leq G} u_k^{-1} D_{k,A} \right)^G$ for $k > 0$, which is an F -isomorphism (Greenlees-Strickland; see also Stapleton).

- ▶ We have identified $\mathrm{spf}(E^0(BA))$ with $\mathrm{Hom}(A^*, \mathbb{G})$.
- ▶ Is there a subscheme of monomorphisms from A^* to \mathbb{G} ?
- ▶ Monomorphisms of schemes are closely related to epimorphisms of commutative rings, which have poor behaviour. Note that $\mathbb{Z} \rightarrow \mathbb{Q}$ and $\mathbb{Z} \rightarrow \mathbb{Z}/n$ are epimorphisms, but behave quite differently.
- ▶ We will define $\mathrm{Level}(A^*, \mathbb{G}) \subseteq \mathrm{Hom}(A^*, \mathbb{G})$ which is motivated by the above; but do not take the analogy too seriously.
- ▶ Write $A = \prod_{i=0}^{s-1} C_{p^{m_i+1}}$, put $D'_A = E^0[[x_i \mid i < s]]$ and $y_i = [p^{m_i}](x_i) \in R'$.
- ▶ Let U be the set of all terms $\sum_{i < s}^F [k_i]_E(y_i)$ with $0 \leq k_i < p$ for all i .
- ▶ Put $g(t) = \prod_{u \in U} (t - u)$ and $[p]_E(t) = q(t)g(t) + r(t)$ with $\deg(r(t)) < s$. Let I be the ideal generated by the coefficients of $r(t)$, and $D_A = D'_A/I$, and $\mathrm{Level}(A^*, \mathbb{G}) = \mathrm{spf}(D_A)$.
- ▶ Schematically:

$$\mathrm{Level}(A^*, \mathbb{G}) = \{ \phi \in \mathrm{Hom}(A^*, \mathbb{G}) \mid \sum_{\alpha \in A^*[\rho]} [\phi(\alpha)] \leq \mathbb{G}[\rho] \}.$$
- ▶ From HKR: $\mathbb{Q} \otimes E^0(BG) = u_0^{-1} E^0(BG) \simeq \left(\prod_{A \leq G} \mathbb{Q} \otimes D_A \right)^G$.
- ▶ There is a similar map $u_k^{-1} E^0(BG)/I_k \rightarrow \left(\prod_{A \leq G} u_k^{-1} D_{k,A} \right)^G$ for $k > 0$, which is an F -isomorphism (Greenlees-Strickland; see also Stapleton).